



Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME							
CENTRE NUMBER					CANDIDATE NUMBER		
CHEMISTRY							9701/31
Paper 3 Advance	ced Practic	al Skills 1				May/Jı	une 2016
							2 hours
Candidates ans	wer on the	Question	Paper.				
Additional Mate	rials: A	s listed in	the Co	onfidential Instructions	S		

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 12 and 13.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

Session	
Laboratory	

For Examiner's Use	
1	
2	
3	
Total	

This document consists of 13 printed pages and 3 blank pages.



1 In this experiment you will determine the identity of the Group 2 metal, **X**, in the carbonate, **X**CO₃. To do this you will react a known mass of **X**CO₃ with **excess** hydrochloric acid, HC*l*, and measure the mass of carbon dioxide that is given off.

FA 1 is XCO₃.

FA 2 is hydrochloric acid, HC*l*.

(a) Method

- Weigh the stoppered tube containing FA 1 and record its mass.
- Use the measuring cylinder to transfer 25 cm³ of **FA 2** into the 250 cm³ beaker.
- Weigh the beaker containing the acid and record the mass.
- Carefully add all the sample of FA 1 to the acid in the beaker.
- Stir the mixture until there is no further reaction.
- Reweigh the beaker and its contents and record the mass.

KEEP THE CONTENTS OF THE BEAKER FOR USE IN QUESTION 2.

- Reweigh the stoppered tube containing any residual FA 1 and record its mass.
- Calculate the mass of FA 1 added to the acid and record this value.
- Calculate the mass of carbon dioxide given off and record this value.

Ι	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of carbon dioxide given off when **X**CO₃ reacted with the acid.

Use the data in the Periodic Table on page 16.

moles of CO ₂	, =	mol
--------------------------	-----	-----

(ii) Write the equation for the reaction of **FA 1**, XCO_3 , with hydrochloric acid, HCl. Include state symbols.

.....

(iii)	Use your answers to (i) and (ii) to calculate the number of moles of XCO_3 that were act to the acid.	lded
	moles of X CO ₃ =	mol
(iv)	Use your answer to (iii) to calculate the relative atomic mass, A_r , of X . Identify X .	
		I
		II
		III
		IV
		V
	A_{r} of $X = \dots$	
	X is	 [5]
	e of the sources of error in this experiment is that it is very difficult to reduce acid spra of the beaker when the metal carbonate is added to the acid.	ying
(i)	Explain what effect this acid spray would have on the value you calculated for the relationic mass, $A_{\rm r}$, of ${\bf X}$.	ative
(ii)	Why is a small amount of acid spray not likely to cause an error in the identification of	 f X ?
()		
(iii)	How could you minimise acid spraying out of the beaker?	
		[3]
	[Total:	: 15]

2 In this experiment you will determine the concentration of the hydrochloric acid, **FA 2**, used in **Question 1**. You will first dilute the reaction mixture that you prepared in **Question 1** and then titrate this diluted solution against sodium hydroxide, NaOH.

$$HCl(aq) + NaOH(aq) \rightarrow NaCl(aq) + H_2O(l)$$

FA 3 is 0.0400 mol dm⁻³ sodium hydroxide, NaOH. methyl orange indicator

(a) Method

Dilution

- Transfer all the reaction mixture that you prepared in **1(a)** from the 250 cm³ beaker to the 250 cm³ volumetric flask.
- Rinse the beaker with a little distilled water and add these washings to the volumetric flask.
- Fill the volumetric flask to the line with distilled water. Stopper the flask and shake it to ensure thorough mixing.
- Label this solution FA 4.

Titration

- Fill the burette with **FA 4**.
- Use a pipette to transfer 25.0 cm³ of **FA 3** into a conical flask.
- Add a few drops of methyl orange.
- Perform a rough titration and record your burette readings in the space below.

The rough	titre i	is	 cm ³

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of FA 4 added
 in each accurate titration.

I II III IV

[4]

(b)		m your accurate titration results, obtain a suitable value for the volume of FA 4 to be used our calculations. Show clearly how you obtained this value.
		25.0 cm ³ of FA 3 required cm ³ of FA 4 . [1]
(c)	Cal	culations
		ow your working and appropriate significant figures in the final answer to each step of your culations.
	(i)	Calculate the number of moles of sodium hydroxide, NaOH, present in 25.0 cm³ of FA 3 .
	(ii)	moles of NaOH = mol Calculate the number of moles of hydrochloric acid, HC $\it l$, present in 250 cm 3 of FA 4 .
((iii)	moles of HCl in 250 cm ³ of $FA 4 = \dots$ mol Use your answers to $1(b)(i)$ and $1(b)(i)$ to calculate the number of moles of HCl that reacted with $FA 1$ in the experiment you carried out in Question 1.
((iv)	moles of HC l that reacted with FA 1 =
		concentration of FA 2 = mol dm ⁻³ [5]

(d)	(i)	One of the sources of error in determining the concentration of FA 2 involves measuring volumes of solutions in both Questions 1 and 2 .
		State which volume of solution that you have measured has the greatest percentage error. How could you have reduced this error?
	(ii)	A student suggested that a greater mass of \mathbf{X} CO $_3$ should be used so that the average titre calculated in $2(\mathbf{b})$ would be a greater volume.
		Explain whether you agree with the student that this would lead to a greater volume for the average titre.
		[2]
		r—1
		[Total: 12]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. Marks are **not** given for chemical equations.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

FA 5 is a mixture of two different salts. Each of these salts contains one cation and one anion from those listed on pages 12 and 13. You will identify the cations and anions present.

(a) (i) Carry out the following test and record your observations.

observations

(ii) Identify one of the cations in	FA 5.
-------------------------------------	-------

One of the cations in **FA 5** is

[2]

(b)	Place the remaining sample of FA 5 in the 100 cm ³ beaker. Half fill the beaker with	distilled
	water and stir until FA 5 has fully dissolved. This may take some time. You will use this	solution
	in the remaining tests.	

(i)	Select reagents to identify the other cation present in FA 5. Carry out tests using these
	reagents and record your results in the space below.
	Identify the cation.

1116 Olliel Calion III I A 3 13	The other	cation	in FA	5 is		
--	-----------	--------	-------	-------------	--	--

(ii) Carry out the following tests and record your observations. Identify one of the anions in **FA 5**.

test	observations
To a 1 cm depth of the solution of FA 5 in a test-tube add aqueous barium chloride or aqueous barium nitrate, then	
add dilute hydrochloric acid.	

One of the anions in **FA 5** is

	9
(iii)	The remaining ion is a halide.
	Select a pair of reagents which can be used to identify the halide present. Carry out a test using these reagents and record your observations below. Suggest the identity of the halide anion present in FA 5 . Explain why this test is not conclusive in this particular case.
	The other anion in FA 5 is

(c)	Suggest the formulae of the two salts that could have been mixed to make FA 5 .	
	and	[1]

[8]

- (d) FA 6 and FA 7 are different organic liquids. Their possible identities are listed below.
 - 2-methylpropan-2-ol
 - propanal
 - propanone

Half fill the 250 cm³ beaker with water and heat to about 50 °C. You will use this as a hot water bath.

Turn off the Bunsen burner.

Carry out the following tests and record your observations.

test	observations
To a 1cm depth of FA 6 in a test-tube, add a few drops of acidified potassium manganate(VII). If no reaction is seen, warm the solution in the hot water bath.	
To a 1cm depth of FA 7 in a test-tube, add a few drops of acidified potassium manganate(VII). If no reaction is seen, warm the solution in the hot water bath.	

Suggest the identity of FA 6 and FA 7 with an explanation.	
FA 6	
FA 7	
	[2]

[Total: 13]

BLANK PAGE

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

ion	reaction with								
ion	NaOH(aq)	NH ₃ (aq)							
aluminium, A <i>l</i> ³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess							
ammonium, NH ₄ +(aq)	no ppt. ammonia produced on heating	_							
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.							
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.							
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess							
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution							
iron(II), Fe²+(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess							
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess							
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess							
manganese(II), Mn²+(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess							
zinc, Zn²+(aq)	white ppt. soluble in excess	white ppt. soluble in excess							

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ -(aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil
nitrite, NO ₂ ⁻ (aq)	NH_3 liberated on heating with $OH^-(aq)$ and Al foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown NO_2 in air)
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl ₂	bleaches damp litmus paper
hydrogen, H ₂	"pops" with a lighted splint
oxygen, O ₂	relights a glowing splint

BLANK PAGE

BLANK PAGE

The Periodic Table of Elements

	18	٥.	<u>Ф</u>	m _i o	0	_o	neon 20.2		_	argon 39.9	9	<u>, -</u>	oton 8	4	Φ.	1.3	9	ے	radon -			
	_	N	エ	heli 4	1	Z	20 ne	-	⋖	arg 39	Ñ		kryr 83	ψ	× 	xer 13	Ø.	~	rad			
	17				6	ш	fluorine 19.0	17	Cl	chlorine 35.5	35	Ŗ	bromine 79.9	53	Н	iodine 126.9	82	Αţ	astatine -			
	16				80	0	oxygen 16.0	16	ഗ	sulfur 32.1	34	Se	selenium 79.0	52	<u>e</u>	tellurium 127.6	84	Ъ	polonium –	116	^	livermorium -
	15				7	z	nitrogen 14.0	15	۵	phosphorus 31.0	33	As	arsenic 74.9	51	Sb	antimony 121.8	83	: <u>.</u>	bismuth 209.0			
	14				9	ပ	carbon 12.0	14	S	silicon 28.1	32	Ge	germanium 72.6	90	Sn	tin 118.7	82	Pb	lead 207.2	114	Εl	flerovium
	13				5	Ф	boron 10.8	13	Ρl	aluminium 27.0	31	Ga	gallium 69.7	49	In	indium 114.8	81	<i>1</i> L	thallium 204.4			
										12	30	Zn	zinc 65.4	48	පි	cadmium 112.4	80	å	mercury 200.6	112	ပ်	copernicium
										7	29	Cn	copper 63.5	47	Ag	silver 107.9	62	Au	gold 197.0	111	Rg	roentgenium -
dn										10	28	z	nickel 58.7	46	Pd	palladium 106.4	78	చ	platinum 195.1	110	Ds	darmstadtium -
Group										6	27	රි	cobalt 58.9	45	뫈	rhodium 102.9	77	٦	iridium 192.2	109	¥	meitnerium -
		_	I	hydrogen 1.0						œ	26	Pe	iron 55.8	44	Ru	ruthenium 101.1	92	Os	osmium 190.2	108	£	hassium
					J					7	25	Mn	manganese 54.9	43	ည	technetium -	75	Re	rhenium 186.2	107	ВР	bohrium
						loc	SS			9	24	ပ်	chromium 52.0	42	Mo	molybdenum 95.9	74	≥	tungsten 183.8	106	Sg	seaborgium -
				Key	atomic number	atomic symbo	name relative atomic mass			2	23	>	vanadium 50.9	41	q	niobium 92.9	73	<u>n</u>	tantalum 180.9	105	<u>ө</u>	dubnium -
					, a	ato	rela			4	22	F	titanium 47.9	40	Zr	zirconium 91.2	72	Ξ	hafnium 178.5	104	¥	rutherfordium -
								_		က	21	Sc	scandium 45.0	39	>	yttrium 88.9	57-71	lanthanoids		89–103	actinoids	
	2				4	Be	beryllium 9.0	12	Mg	magnesium 24.3	20	Ca	calcium 40.1	38	Š	strontium 87.6	56	Ba	barium 137.3	88	Ra	radium
	_				3	:=	lithium 6.9	1	Na	sodium 23.0	19	×	potassium 39.1	37	Rb	rubidium 85.5	55	S	caesium 132.9	87	Ļ	francium -

r ₇ Lu	lutetium 175.0	103	۲	lawrencium	ı
oz Yb	ytterbium 173.1	102	Š	nobelium	1
e9 Tm	thulium 168.9	101	Md	mendelevium	1
₆₈	erbium 167.3	100	Fn	ferminm	ı
67 Ho	holmium 164.9	66	Es	einsteinium	1
66 Dy	dysprosium 162.5	86	ర్	californium	1
e5 Tb	terbium 158.9	26	Ř	berkelium	ı
² DD	gadolinium 157.3	96	CB	curium	ı
63 Eu	europium 152.0	98	Am	americium	ı
62 Sm	samarium 150.4	94	Pn	plutonium	ı
Pm	promethium -	93	ď	neptunium	ı
 9 V	neodymium 144.4	92	\supset	uranium	238.0
eg Ā	praseodymium 140.9	91	Ра	protactinium	231.0
Çe Ce	cerium 140.1	06	Ļ	thorium	232.0
57 La	lanthanum 138.9	89	Ac	actinium	ı

lanthanoids

actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.