



Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

AS & A Level			
CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			9701/34
Paper 3 Advance	ed Practical Skills 2	Oct	tober/November 2016
			2 hours
Candidates answe	er on the Question Paper.		
Additional Materia	als: As listed in the Confidential In	structions	
READ THESE IN	STRUCTIONS FIRST		
Give details of the Write in dark blue You may use an I Do not use staple	HB pencil for any diagrams or graphs. es, paper clips, glue or correction fluid.	ere appropriate, in the boxes pr	rovided.
DO NOT WRITE	IN ANY BARCODES.		

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11. A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document consists of 12 printed pages.



You will find the relative atomic mass, A_r , of magnesium by measuring the volume of hydrogen produced when a known mass of metal reacts with an excess of acid.


$$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$$

FB 1 is 1.00 mol dm⁻³ hydrochloric acid, HC*l*. **FB 2** is magnesium, Mg.

(a) Method

Read through the whole method before starting any practical work.

- Fill the tub with water to a depth of about 5 cm.
- Weigh the magnesium, FB 2, and note its mass below. If you are using a balance reading
 to 1 decimal place and the reading with the magnesium is zero, you should record this
 value.
- Fill the burette to about the 20 cm³ mark with hydrochloric acid, **FB 1**.
- Add distilled water to reach the 0 cm³ mark on the burette.
- Bend the magnesium strip into a U-shape.
- Place the magnesium in the burette so that it is above the liquid and friction holds it in position. Use a glass rod to push the magnesium about 2 cm into the burette.
- Hold a piece of paper towel over the open end of the burette, invert the burette and immediately place it in the tub of water. Remove the paper towel and clamp the burette as shown in the diagram.
- The liquid level should now be on the scale of the burette. If it is not, open the tap for a
 moment to allow the level to drop.



- Record the initial reading on the burette. Remember that the scale is now upside down.
- Leave the apparatus so that the acid from the burette diffuses around the magnesium and reacts.
- You should start Question 2 or Question 3 while waiting for the reaction to complete.
- When all the magnesium has reacted, note and record the final reading on the burette.
- Calculate the volume of hydrogen produced.

Results

((b)	Cal	lcu	latio	ns
۱		, va	ıvu	ıaıı	7113

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

	(i)	Calculate the number of moles of hydrogen produced. (Assume that 1 mole of gas occupies 24.0 dm³ under these conditions.)
	(ii)	moles of H_2 =
(c)	(i)	$A_{\rm r}$ of Mg =[2] Calculate the percentage error in the mass and volume readings in this experiment.
	(ii)	Suggest a change that could be made to reduce the greater error calculated in (i).
(d)		[3] at would be the effect on the value of the A_r of magnesium calculated if the temperature of room was much lower than that for your experiment? Explain your answer.

[Total: 10]

2 In **Question 1** you calculated the relative atomic mass, *A*_r, of magnesium by measuring the volume of hydrogen produced. The relative atomic mass can also be determined by investigating how much of the hydrochloric acid reacted with the magnesium.

The experiment described in **Question 1** was repeated, this time using 0.21 g of magnesium ribbon and 30.0 cm³ of 1.00 mol dm⁻³ hydrochloric acid. All the solution left in the burette and tub was kept and water added to make the total volume 250 cm³. This solution was labelled **FB 3**.

You will titrate **FB 3** using a known concentration of aqueous sodium carbonate to determine how much hydrochloric acid was left over after the reaction with magnesium.

$$Na_2CO_3(aq) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l) + CO_2(g)$$

FB 3 is the solution of hydrochloric acid described above.

FB 4 is aqueous sodium carbonate containing $2.64\,\mathrm{g\,dm^{-3}\,Na_2CO_3}$. bromophenol blue indicator

(a) Method

- Fill the burette with **FB 3**.
- Pipette 25.0 cm³ of **FB 4** into a conical flask.
- Add about 10 drops of bromophenol blue indicator.
- Perform a rough titration and record your burette readings in the space below.

The rough titi	e is	cm ³
----------------	------	-----------------

- Carry out as many accurate titrations as you consider necessary to obtain consistent results.
- Record, in a suitable form below, all of your burette readings and the volume of FB 3 added in each accurate titration.
- Make certain any recorded results show the precision of your practical work.

I II III IV V VI VII VIII

[7]

(b)		om your accurate titration results, obtain a suitable value for the volume of FB 3 to be used your calculations. Show clearly how you obtained this value.
		25.0 cm ³ of FB 4 required cm ³ of FB 3 . [1]
(c)	Cal	lculations
		ow your working and appropriate significant figures in the final answer to each step of your culations.
	(i)	Use the information on page 4 and the Periodic Table on page 12 to calculate the number of moles of sodium carbonate in the 25.0 cm ³ of FB 4 used in each titration.
		moles of Na ₂ CO ₃ = mol
	(ii)	Use your answer to (i) to calculate the number of moles of hydrochloric acid present in the volume of FB 3 recorded in (b).
		$Na_2CO_3(aq) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(I) + CO_2(g)$
		moles of HC1 present = mol
((iii)	Use your answer to (ii) to calculate the number of moles of hydrochloric acid present in 250 cm³ of FB 3 .
		moles of HC l present in 250 cm 3 = mol

(iv)	Use the information on page 4 to calculate the number of moles of hydrochloric acid added to the magnesium.
	moles of HCl added = mol
(v)	Calculate the number of moles of hydrochloric acid that reacted with the magnesium.
	moles of HC <i>l</i> that reacted with the magnesium = mol
(vi)	Use your answer to (v) and the mass of magnesium used to calculate the relative atomic mass, A_r , of magnesium.
	$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$
	A_{r} of Mg =[5]
	olution of sodium hydroxide was prepared at the same concentration, in moldm ⁻³ , as FB 4 . tudent repeated the titration but replaced FB 4 with this solution of sodium hydroxide.
(i)	Explain the effect that replacing FB 4 with this solution of sodium hydroxide would have on the volume of acid, FB 3 , needed for the titration.
(ii)	If the sodium hydroxide had been stored for a long time it would not be suitable for use to find the concentration of the acid.
	Suggest why storage for a long time would make the sodium hydroxide unsuitable.
	[2]
	[Tatal: 45]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) (i) Half fill the 250 cm³ beaker with water. Heat the water to about 80 °C and then turn off the Bunsen burner. This is the hot water bath needed in the following tests.

To a 3–4 cm depth of aqueous silver nitrate in a test-tube, add a few drops of aqueous sodium hydroxide to give a grey/brown precipitate. Then add aqueous ammonia dropwise until the precipitate **just** disappears. This solution is Tollens' reagent and is needed in a following test.

FB 5, FB 6 and FB 7 are each known to be one of ethanol, propanal and propanone.

observations

Carry out the following tests and complete the table.

test

test				
		FB 5	FB 6	FB 7
To a 1 cm depth in a test-tube, add a few drops of acidified potassium manganate(VII) and place in the hot water bath.				
To a 0.5 cm depth in a test-tube, add a 1 cm depth of aqueous potassium iodide and a 1 cm depth of sodium chlorate(I). (This gives the same result as adding iodine and alkali.)				
To a few drops in a test-tube, add a 1 cm depth of Tollens' reagent and place in the hot water bath. Leave for several minutes.				
(ii)	Use these observation	ns to identify the unkno	wn compounds.	
	FB 5 is			
	FB 6 is			
	FB 7 is			
(iii)	Choose another reage different result for ethat	ent that would give a sir anol.	milar result for propana	l and propanone but a
	Do not carry out this	test.		
	reagent			
	result for propanal and	d propanone		
	result for ethanol			
(iv)	Choose another reage different result for prop	ent that would give a si panal.	milar result for ethanol	and propanone but a
	Do not carry out this	test.		
	reagent			
	result for ethanol and	propanone		
	result for propanal			[8]
				F-1

(b)		8 contains one cation and one anion from those listed on pages 10 and 11. a are provided with solid FB 8 and an aqueous solution of FB 8.
	(i)	To a 1cm depth of aqueous FB 8 in a test-tube add a 1cm depth of aqueous sodium hydroxide.
		Keep the test-tube and contents for test (ii).
		observation
	(ii)	Transfer the contents of the test-tube from test (i) into a boiling tube and heat gently and carefully.
		Allow to cool and keep the boiling tube and contents for test (iii).
		observation
((iii)	Transfer a 1 cm depth of the mixture from test (ii) into a boiling tube and add a 2 cm depth of dilute hydrochloric acid. Heat gently and carefully.
		observation
		Allow to cool and keep the boiling tube and contents for test (iv).
((iv)	To the boiling tube from test (iii) add a piece of aluminium foil. Leave the boiling tube to stand.
		observation
	(v)	Place a small spatula measure of solid FB 8 in a hard-glass test-tube and heat it gently at first and then more strongly.
		Identify two gases, other than water vapour, that are produced and give your evidence.
		identity
		evidence
		identity
		evidence
((vi)	From your observations in (i) to (v), write the formula of FB 8.
(\	vii)	Write the ionic equation for the reaction that is occurring in test (i). Include state symbols.
		[7]
		L'1

[Total: 15]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

ion	reaction with		
ion	NaOH(aq)	NH ₃ (aq)	
aluminium, Al³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess	
ammonium, NH₄⁺(aq)	no ppt. ammonia produced on heating	_	
barium, Ba ²⁺ (aq)	no ppt. (if reagents are pure)	no ppt.	
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca2+(aq)]	no ppt.	
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess	
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution	
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess	
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess	
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess	
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess	
zinc, Zn²+(aq)	white ppt. soluble in excess	white ppt. soluble in excess	

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ -(aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil
nitrite, NO ₂ ⁻ (aq)	NH_3 liberated on heating with $OH^-(aq)$ and Al foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown NO_2 in air)
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl ₂	bleaches damp litmus paper
hydrogen, H ₂	"pops" with a lighted splint
oxygen, O ₂	relights a glowing splint

The Periodic Table of Elements

	~		(I)	E ~		(I)	c 0		<u> </u>	<u>د</u> ه		<u> </u>	5 &		a)	ار اد دن		_	5				
	18	2	Ĭ	heliu 4.0	10	ž	20.5	18	₹	argon 39.9	36	Ž	krypt 83.5	22	×	xenc 131	86	Ř	rado				
	17				6	Щ	fluorine 19.0	17	Cl	chlorine 35.5	35	Ā	bromine 79.9	53	П	iodine 126.9	85 At astatine						
	16				8	0	oxygen 16.0	16	S	sulfur 32.1	34	Se	selenium 79.0	52	<u>e</u>	tellurium 127.6	84	Ро	polonium –	116	_	livermorium	ı
	15				7	z	nitrogen 14.0	15	۵	phosphorus 31.0	33	As	arsenic 74.9	51	Sb	antimony 121.8	83	<u>B</u>	bismuth 209.0				
	14				9	O	carbon 12.0	14	S	silicon 28.1	32	Ge	germanium 72.6	20	S	tin 118.7	82	Ър	lead 207.2	114	Εl	flerovium	ı
	13				2	В	boron 10.8	13	Ρl	aluminium 27.0	31	Ga	gallium 69.7	49	In	indium 114.8	81	l1	thallium 204.4				
										12	30	Zu	zinc 65.4	48	පි	cadmium 112.4	80	Р	mercury 200.6	112	ပ်	copernicium	1
										7	29	Cu	copper 63.5	47	Ag	silver 107.9	62	Αn	gold 197.0	111	Rg	roentgenium	1
dn										10	28	Ē	nickel 58.7	46	Pd	palladium 106.4	78	置	platinum 195.1	110	Ds	darmstadtium	ı
Group										6	27	ပိ	cobalt 58.9	45	뫈	rhodium 102.9	77	'n	iridium 192.2	109	¥		
		-	I	hydrogen 1.0						80	26	Ь	iron 55.8	44	Ru	ruthenium 101.1	92	SO	osmium 190.2	108	Ϋ́	hassium	ı
					J					7	25	M	manganese 54.9	43	ပ	technetium -	75	Re	rhenium 186.2	107	В	pohrium	ı
						00	SS			9	24	ပ်	chromium 52.0	42	Мо	molybdenum 95.9	74	>	tungsten 183.8	106	Sg	seaborgium	ı
				Key	atomic number	atomic symbo	name relative atomic mass			2	23	>	vanadium 50.9	41	g	niobium 92.9	73	ц	tantalum 180.9	105	9	dubnium	-
					ъ	ator	relat			4	22	F	titanium 47.9	40	Zr	zirconium 91.2	72	Ξ	hafnium 178.5	104	꿒	rutherfordium	
								1		က	21	Sc	scandium 45.0	39	>	yttrium 88.9	57-71	lanthanoids		89-103	actinoids		
	2				4	Be	beryllium 9.0	12	Mg	magnesium 24.3	20	Ca	calcium 40.1	88	ഗ്	strontium 87.6	26	Ba	barium 137.3	88	Ra	radium	
	_				е	=	lithium 6.9	1	Na	sodium 23.0	19	×	potassium 39.1	37	Rb	rubidium 85.5	55	S	caesium 132.9	87	ъ̈́	francium	1

71 Lu	175.0	Ļ	lawrencium	ı
70 Yb	173.1	8 N	nobelium	ı
69 Tm	168.9	Md	mendelevium	ı
68 Fr erbium	167.3	Fm	ferminm	ı
67 Ho	164.9	Es	einsteinium	ı
66 Dy dysprosium	162.5	Ç	californium	ı
65 Tb	158.9	Ř	berkelium	ı
Gd gadolinium	157.3	Cm	curium	ı
63 EU europium	152.0	Am	americium	ı
62 Sm	150.4	Pn	plutonium	ı
61 Pm promethium	- 83	ď	neptunium	ı
60 Nd neodymium	144.4	⊃	uranium	238.0
Pr praseodymium	140.9	Ра	protactinium	231.0
Ce certum	140.1	H	thorium	232.0
57 La	138.9	Ac	actinium	ı

lanthanoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.