



Cambridge Assessment International Education

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		
CHEMISTRY			0620/62	
Paper 6 Alternative to Practical			February/March 2019	
			1 hour	
Candidates ans	Candidates answer on the Question Paper.			

READ THESE INSTRUCTIONS FIRST

No Additional Materials are required.

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate. This document consists of 9 printed pages and 3 blank pages.



1 A chemist heated solid zinc chloride until it became molten. The apparatus shown was then used to pass electricity through the molten zinc chloride using inert electrodes.



A silver-coloured solid was formed at the negative electrode (cathode).

(a)	Name the process of breaking down a substance using electricity.	
		[1]
(b)	A Bunsen burner was used to heat the zinc chloride.	
	Describe how a Bunsen burner is adjusted to give a very hot flame.	
		[1]
(c)	Suggest and explain the expected observation at the positive electrode (anode).	
		[4]
(d)	Suggest why iron electrodes cannot be used in this experiment.	

(e)	(i)	What difference would the chemist observe at the negative electrode if aquezinc chloride were used, rather than molten zinc chloride? Explain you answer.	ous
		difference	
		explanation	
			[2]
	(ii)	When electricity is used to break down concentrated aqueous zinc chloride, chloring produced at the positive electrode.	e is
		Describe a test for chlorine.	
		test	
		observations	[2]
(f)	The	e bottle of zinc chloride is labelled <i>corrosive</i> .	
	Sta	te one safety precaution that should be taken when using zinc chloride.	
			[1]
		[Total:	101

2 A student investigated the reaction between two different solutions, **A** and **B**, of aqueous potassium manganate(VII) and solution **C**.

Three experiments were done.

Experiment 1

- A burette was filled with solution A. The initial burette reading was recorded.
- A measuring cylinder was used to pour 25 cm³ of solution **C** into a conical flask.
- Solution A was added to the conical flask until the mixture just turned pink. The final burette reading was recorded.
- About 2 cm³ of the contents of the conical flask was poured into a test-tube to use in Experiment 3.
- The rest of the contents of the conical flask was poured away. The conical flask was rinsed with distilled water.
- (a) Use the burette diagrams to record the burette readings in the table and complete the table.



initial burette reading

final burette reading

	Experiment 1
final burette reading/cm ³	
initial burette reading/cm ³	
volume used/cm ³	

[2]

Experiment 2

- The contents of the burette used in Experiment 1 were poured away and the burette was rinsed with distilled water.
- The burette was then rinsed with solution **B**.
- Experiment 1 was repeated using solution B instead of solution A.
- (b) Use the burette diagrams to record the burette readings in the table and complete the table.



initial burette reading

final burette reading

	Experiment 2
final burette reading/cm ³	
initial burette reading/cm³	
volume used/cm ³	

[2]

(c) (i) Which solution of potassium manganate(VII), solution **A** or solution **B**, is the more concentrated?

Explain your answer.

[2]

(ii) How many times more concentrated is this solution of potassium manganate(VII)?

......[1]

(d) (i) Predict the volume of solution B that would be used if Experiment 2 were rep 50 cm³ of solution C . Explain your answer.			oe used if Experiment 2 were repeat	ed using	
	(ii)	Suggest a practical problem that using 50 cr			
		problem be solved?			
				[2]	
(e)		e one advantage and one disadvantage of usi solution C .	ng a measuring cylinder rather than	a pipette	
	adv	antage of using a measuring cylinder			
	disa	disadvantage of using a measuring cylinder			
				[2]	
Exp	perim	nent 3			
The	e resi	ults from Experiment 3 are shown in the table			
		tests	observations		
		queous sodium hydroxide was added to cout 2 cm³ of solution C .	green precipitate formed		
		queous sodium hydroxide was added to the eaction mixture saved from Experiment 1.	red-brown precipitate formed		
(f)	Wh	at conclusions can be drawn about solution C	from Experiment 3?		
				[2]	

0620/62/F/M/19

© UCLES 2019

[Total: 15]

3		substances, solution D and solid E , were analysed. Solution D was dilute sulfuric acid. ts were done on the substances.
	tes	ts on solution D
	Cor	mplete the expected observations.
	Sol	ution D was divided into four equal portions in four test-tubes.
	(a)	The pH of the first portion of solution D was tested.
		pH =[1]
	(b)	A strip of magnesium ribbon was added to the second portion of solution D . The gas produced was tested.
		observations
		[3]
	(c)	Dilute nitric acid and aqueous silver nitrate were added to the third portion of solution D .
		observations[1]

(d) Dilute nitric acid and aqueous barium nitrate were added to the fourth portion of solution **D**.

tests on solid E

Some of the tests and observations are shown.

tests on solid E	observations
The appearance of solid E was studied.	white solid
test 1	
Solid E was heated gently and then more strongly.	white solid residue
Distilled water was added to the residue and the pH of the mixture was tested.	pH = 10
test 2	
Dilute hydrochloric acid was added to solid E .	rapid effervescence
The gas produced was tested.	limewater turned milky
Distilled water was added to the solution and the mixture was shaken.	
An excess of aqueous sodium hydroxide was added to the mixture.	white precipitate formed which was insoluble in excess

(e)	Identify the gas produced in test 2.	
		[1]
(f)	What conclusions can you draw about solid E ?	
		[2]
	[Total	l: 9]

4 The rate of reaction between magnesium and dilute hydrochloric acid can be followed by measuring the volume of hydrogen produced.

Plan an experiment to investigate the effect of decreasing the temperature on the rate of this reaction by measuring the volume of hydrogen produced.

You are provided with magnesium ribbon, dilute hydrochloric acid and common laboratory apparatus.

You are advised to draw a labelled diagram of the apparatus you would use in the space provided.

 [6]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.