MARK SCHEME for the May/June 2015 series

0620 CHEMISTRY

0620/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.



Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Abbreviations used in the Mark Scheme

- •; separates marking points
- / separates alternatives within a marking point
- OR gives alternative marking point
- R reject
- I ignore mark as if this material was not present
- A accept (a less than ideal answer which should be marked correct)
- COND indicates mark is conditional on previous marking point
- owtte or words to that effect (accept other ways of expressing the same idea)
- max indicates the maximum number of marks that can be awarded
- ecf credit a correct statement that follows a previous wrong response
- () the word / phrase in brackets is not required, but sets the context
- ORA or reverse argument

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
1(a)	Any two fossil fuels from: crude oil/petroleum; natural gas/methane; petrol/gasoline; kerosene/paraffin; diesel (oil)/gas oil; fuel oil; refinery gas/LPG; propane; butane;	2	I ethane/oil/naphtha/coal/gas R coke/bitumen/lubricating oil/wood
1(b)	hydrogen, oxygen, nitrogen; All three for 2 marks two for 1 mark	2	A H, O, N I H ₂ , O ₂ , N ₂
1(c)(i)	M1 oxygen and nitrogen (from air) react; M2 oxides of nitrogen OR nitrogen oxide(s) are formed;		 A nitrogen combust for M1 R M1 if oxygen or nitrogen originate from the fuel A named oxide of nitrogen e.g. nitrogen dioxide A correct formulae A NO_x
	M3 nitrogen oxides formed react with water (to form acid);	3	

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
1(c)(ii)	Any two from:		R 'global warming/greenhouse effect'
	M1 lowers pH or acidifies lakes/rivers or kills fish;		${\bf R}$ 'increases pH of lakes so kills fish' for M1
	M2 changes composition of soils or reduces fertility of soil or reduces crop yields deforestation or kills crops/trees/plants/leaves;		A removes nutrients/leaches the soil
	M3 attacks (limestone) buildings or statues;		A alternative words for 'attacks' e.g. damages/reacts with/corrode/erode for M3 and M4
	M4 attacks metal (structures)/bridges;	3	I rusting but A 'enhances rusting' for M4 I toxicity to humans
1(d)	Any three from: M1 wood burns to produce (less) carbon dioxide; M2 trees (wood) take in carbon dioxide; M3 by photosynthesis; M4 wood is carbon neutral fuel;	3	

[Page 5	Mark Scheme	Syllabus	Paper
		Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
2(a)	M1 <i>Forming an oxide</i> (all) elements or (all) impurities become oxides;		 (All) elements or (all) impurities react with oxygen A M1 for any one element becoming an oxide
	M2 Gaseous oxides carbon dioxide or sulfur (di)oxide escape/are removed as gases;		A formulae/carbon monoxide A oxides of sulfur/carbon I sulfur trioxide
	M3 Acidic oxides silicon(IV) oxide or phosphorus(III/V) oxide react/are neutralised by calcium oxide/lime;		A silicon (di)oxide for silicon(IV) oxide A phosphorus (tri/pent)oxide for phosphorus(III/V) oxide
	M4 Equation mark any one of the following equations $S + O_2 \rightarrow SO_2;$ $C + O_2 \rightarrow CO_2 \text{ or } 2C + O_2 \rightarrow 2CO;$ $Si + O_2 \rightarrow SiO_2;$ $4P + 5O_2 \rightarrow 2P_2O_5 \text{ or } P_4 + 5O_2 \rightarrow 2P_2O_5;$ $4P + 3O_2 \rightarrow 2P_2O_3 \text{ or } P_4 + 3O_2 \rightarrow 2P_2O_3;$		A multiples I state symbols I unbalanced equations R other combustion equations with incorrect species
	M5 Word equation mark any one of the following word equations calcium oxide + silicon(IV) oxide \rightarrow calcium silicate; calcium oxide + phosphorus(III/V) oxide \rightarrow calcium phosphate;	5	A calcium oxide + silicon(IV) oxide \rightarrow slag A correct symbol equation for M5 but R other equations with incorrect species used as M5

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
2(b)(i)	Any one from: (making) car (bodies); machinery; chains; pylons; white goods; nails; screws; as a building material; sheds/roofs; reinforcing concrete;	1	A bridges A tools I cutlery
2(b)(ii)	Any one from: knives; drills; railway tracks; machine/cutting tools/hammers; razor blades; chisels;	1	I cutlery items I bridges
2(b)(iii)	M1 atoms or cations or (positive) ions or metal ions; M2 arranged in a lattice or in layers or in rows or in a regular structure; M3 rows or layers slide over one another;	3	 I (sea of) electrons R protons or nuclei for M1 A M2 non-directional forces A ECF on particle named in M1 for M3 I 'atoms' slide over one another
2(b)(iv)	M1 carbon atoms or particles in structure different size (to cations); M2 so reduce moving or interrupt movement;	2	 R ions and molecules for M1 A M2 for prevents sliding A M2 for 'stops' sliding
3(a)(i)	Zn to Zn ²⁺ ; because electron loss;	2	A because oxidation number has increased for M2

Page	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
3(a)(ii)	(2)H ⁺ or 'hydrogen ion(s)';		R H ₂ or 'hydrogen'
	it accepts electrons or takes electrons (from zinc atoms);	2	 A because it is reduced or because it decreases in oxidation number A it causes zinc to lose electrons
3(b)(i)	zinc displaces copper or zinc more reactive than copper;		A copper less reactive than zinc I zinc reacts with copper ions or with Cu ²⁺ or with copper chloride I zinc reacts with copper I Cu ²⁺ ions are reduced
	Zn + CuC l_2 → ZnC l_2 + Cu OR Zn + Cu ²⁺ → Cu + Zn ²⁺ ;	2	A multiples I state symbols
3(b)(ii)	steeper (line) or higher gradient; (means an) increased rate;		A less time to complete the reaction/same amount of gas in less time/faster reaction/ more gas in the same time period
	but the same (final) volume;	3	 A same volume of hydrogen produced A 'amount' for volume A no extra gas is made
3(c)	M1 less steep (line) or lower gradient;		A alternative phrases e.g. 'shallower'
	M2 (because of) decreased rate;		 A more time to complete the reaction A same amount of gas in more time A slower rate or slower reaction
	M3 ethanoic is a weak(er) acid;		ORA
	M4 only partially ionised or dissociated OR lower concentration of hydrogen ions;	4	A not fully dissociated or ionised A ionises less (than HC <i>l</i>) I less hydrogen ions

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
3(d)	M1 moles of HC $l = 0.1$ (mol);		
	M2 moles of $Zn = 0.05 (mol);$		A ECF for M1 \times $\frac{1}{2}$
	mass of zinc = 3.25g;	3	A ECF for M2 \times 65 Unit required for M3
4(a)(i)	<i>Any three from:</i> same general formula; contain the same functional group; consecutive members differ by CH ₂ ; common methods of preparation; same or similar chemical properties; physical properties vary in a predictable manner / show trends / show a gradual change / an example of a physical variation e.g. mpt, bpt volatility viscosity;	3	I different physical properties / physical properties change / an unqualified or slight change R same or similar physical properties
4(a)(ii)	propanol/propan-1-ol/propan-2-ol;	1	
4(a)(iii)	if molecular formula is given as $C_{10}H_{22}O$ award 2 marks		A C ₁₀ H ₂₁ OH for two marks
	if not, look for evidence of some correct working for one mark $158 - 17 = 141$ OR $12n + 2n + 1 = 141$ OR $n = 10$	2	A (10 × 12) + (22 × 1) + 16 = 158 for one (working) mark
4(b)	they have the same molecular formula ($C_4H_{10}O$);		A same number of each type of atom I same number of atoms
	different structures;	2	A different structural formula or different arrangement of atoms

Page 9	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
4(c)(i)	M1 butene or but-1-ene;		M1 and M2 are independent A but-2-ene for M1
	M2 structural formula of but-1-ene;	2	Minimum acceptable structure is $CH_3CH_2CH=CH_2$ Double bond must be shown R structure of but-2-ene for M2
4(c)(ii)	butyl ethanoate;	1	A butanyl R ethenoate and ethanoic
4(c)(iii)	butanoic acid; structural formula of butanoic acid;	2	A butyric acid Minimum acceptable structure is CH ₃ CH ₂ CH ₂ CO ₂ H A CH ₃ CH ₂ CH ₂ COOH with C–HO connectivity in acid group
5(a)	M1 add chlorine to (potassium) iodide solution;		Solution must be implied for M1 A any soluble iodide solution
	M2 red/brown/yellow/orange (solution) is formed;		A black (ppt or solid)
	$\begin{array}{rcl} M3 \\ Cl_2 &+& 2KI \rightarrow & 2KCl &+& I_2 \\ Cl_2 &+& 2I^- \rightarrow & 2Cl^- &+& I_2 \end{array}; \end{array}$	3	A multiples I state symbols but KI(aq) would allow the solution aspect of mark in M1
5(b)	M1 (0.013 moles of I and 0.065 moles of F atoms gives a) ratio 1:5;		Award 2 marks for IF_5
	Formula = IF ₅ ;	2	A one mark for I_5F (as ratio is inverted) A one mark for IFl_5 or I_5Fl

Γ	Page 10	Mark Scheme	Syllabus	Paper
		Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
5(c)(i)	example of a reversible reaction including attempts at removing/adding waters of crystallisation OR example of a reaction which under closed conditions would be reversible;	1	 A written description of the reaction e.g. 'Haber process' unless equation is attempted in which case ignore written description A word equations / unbalanced equations A equations without equilibrium arrows I descriptions of physical changes
5(c)(ii)	Any two from: (a reaction) M1 which can take place in both directions OR which can be approached from both directions;		I reference to 'closed system' A 'a reaction which can go forwards and backwards' for M1 I 'a reaction with an equilibrium arrow' or with '≓' for M1
	M2 in which concentrations/macroscopic properties do not change (with time);		R concentrations (of reactants and products) are the same
	M3 the two reaction rates are equal;	2	
5(d)	M1 equilibrium goes to LHS OR equilibrium goes to reactants side;		 A reaction goes to LHS but R 'equilibrium goes to LHS and to products side' A backward reaction is favoured I less yield or less products
	M2 because the concentration of chlorine decreases;	2	 A 'reactant' for 'chlorine' but not reactants A to replace missing chlorine

Page '	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
5(e)	M1 equilibrium goes to RHS OR equilibrium goes to products side;		 A reaction goes to RHS but R 'equilibrium goes to RHS and to reactants side' A forward reaction is favoured I more yield or more products
	M2 exothermic reactions are favoured by low temperatures;		A for M1 and M2 'decreasing temperature makes the equilibrium go to RHS'
	M3 the forward reaction is exothermic;	3	A backward reaction is endothermic

Page 12	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
6(a)(i)	M1 proton acceptor;		A alternative words to 'acceptor' e.g. 'receiver' I references to pH
	M2 does not accept (protons) readily OR less able to accept protons (than strong bases);	2	A 'hydrogen ion' or 'H ⁺ ' for proton I accepts fewer/less protons
6(a)(ii)	M1 same <u>concentration</u> of both bases;		
	M2 measure their pH;		 A suitable method e.g. universal indicator or pH paper or pH meter I litmus or methyl orange or phenolphthalein I titration methods for M2 and M3
	M3 the higher pH is the stronger base;	3	 A suitable colours of both weak strong bases e.g. ethylamine is (greeny)blue, NaOH is darker blue/purple A alternative methods for M2 and M3 e.g. measure conductivity (M2) and higher conductivity is the stronger base (M3) e.g.add aluminium/Al (M2) and stronger base gives faster rate of effervescence/more fizzing/more bubbling (M3)
6(b)(i)	$\begin{array}{rcl} 2CH_{3}CH_{2}NH_{2} + & H_{2}SO_{4} \rightarrow & (CH_{3}CH_{2}NH_{3})_{2}SO_{4} \\ \text{species;} \\ \text{balancing;} \end{array}$		 A multiples I state symbols A one mark for correct product
	the salt is ethylammonium sulfate;	3	A close spellings A diethylammonium sulfate

Page 13	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2015	0620	31

Question	Answer	Marks	Guidance
6(b)(ii)	sodium hydroxide / calcium hydroxide / NaOH / Ca(OH) ₂ ;	1	A any Group 1 or Group 2 hydroxide or oxide
6(c)(i)	<i>Any two from:</i> (particles move in) random motion;		
	(particles) collide;		A alternative phrases for collide
	(particles) move from a region of high concentration to low concentration;	2	A down a concentration gradient
6(c)(ii)	C; M2 it has a lower (relative) molecular mass (than HBr); M3 ethylamine diffuses faster (than HBr);	3	 A ethylamine is less dense A ethylamine is a lighter molecule but I 'ethylamine is lighter' I ethylamine is a smaller molecule A ethylamine molecules or particles move faster A ECF for M2 and M3 if A is given e.g. HBr diffuses faster for M3 because it is a lighter molecule for M2 A ECF for M2 if B is given e.g. they diffuse at same rate for M3 because molecules weigh the same for M2