

Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER	CANDID/ NUMBEF	
CHEMISTRY		0620/51
Paper 5 Practical Test May/June 2		
		1 hour 15 minutes
Candidates ans	swer on the Question Paper.	
Additional Mate	erials: As listed in the Confidential Instructions	
READ THESE	INSTRUCTIONS FIRST	

Write your Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use an HB pencil for any diagrams or graphs.Do not use staples, paper clips, glue or correction fluid.DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units. Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

Total

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 6 printed pages and 2 blank pages.



You are going to investigate what happens when dilute hydrochloric acid reacts with an aqueous solution \mathbf{R} , containing two different substances \mathbf{S} and \mathbf{T} .

Read all the instructions below carefully before starting the experiments.

Instructions

1

You are going to carry out three experiments.

(a) Experiment 1

Using a measuring cylinder, pour 25 cm^3 of solution **R** into the conical flask. Add five drops of the phenolphthalein indicator to the flask.

Fill the burette with hydrochloric acid to the 0.0 cm^3 mark. Using the burette, add hydrochloric acid to solution **R** and shake the flask. Continue to add hydrochloric acid to the flask until the colour just disappears.

Record the volume of hydrochloric acid added when the indicator changes colour. **Keep the mixture** in the flask for Experiment 2.

	burette readings	
final volume/cm ³		
initial volume/cm3		
difference / cm ³		

(b) Experiment 2

Add four drops of methyl orange indicator to the mixture in the flask from Experiment 1. Check that the burette reading is the same as your final reading in Experiment 1. This is the initial volume reading for Experiment 2.

Using the burette, add hydrochloric acid to the mixture in the flask and shake the mixture. Record the volume of hydrochloric acid added when the indicator just changes colour.

	burette readings	
final volume/cm ³		
initial volume/cm ³		
difference / cm ³		

[4]

[3]

(c) Experiment 3

Pour about 5 cm^3 of solution **R** into a test-tube. Using a teat pipette, add hydrochloric acid to the solution. Note any observations.

......[1]

(d) (i)	When phenolphthalein indicator was used in Experiment 1 the colour changed
	from [2]
(ii)	What was the colour of the indicator at the end-point in Experiment 2?
(e) (i)	What type of substances are S and T ? [1]
(ii)	What conclusion can you draw from Experiment 3?
	[1]
	e volume of hydrochloric acid added in Experiment 1 reacted with all of substance S and half substance T .
	e volume of hydrochloric acid in Experiment 2 reacted with half of substance T .
(i)	Work out the volume of hydrochloric acid which reacted with substance ${f S}$.
(ii)	Work out the volume of hydrochloric acid which reacted with substance T .
(iii)	Compare the volumes of hydrochloric acid which reacted with substances ${f S}$ and ${f T}$.
(g) (i)	Predict the volume of hydrochloric acid which would be added in Experiments 1 and 2 if the experiments were repeated using 100cm^3 of solution R . Explain your answer.
	Experiment 1
	Experiment 2
	Explanation[3]
(ii)	Suggest a practical problem that would occur when carrying out these repeat experiments and how you could solve this problem.
	[Total: 22]

2 You are provided with solid **U**, which is a metal salt. Carry out the following tests on **U**, recording all of your observations in the table. Conclusions must **not** be written in the table.

	tests	observations
tests on solid U		
(a)	Describe the appearance of solid U .	[2]
(b)	Place half of solid U in a test-tube. Heat the test-tube gently at first and then more strongly.	[2]
(c)	Add the rest of solid U to about 8 cm ³ of distilled water in a test-tube. Stopper the test-tube and shake the contents until dissolved. Divide the solution into four equal portions in separate test-tubes and carry out the following tests.	
	Add several drops of aqueous sodium hydroxide to the first portion of the solution and shake the test-tube. Then add about 1 cm ³ of hydrogen peroxide solution to the mixture.	
	Test the gas given off.	[5]
(d)	Add excess aqueous ammonia to the second portion of the solution.	[1]
(e)	Add about 1 cm ³ of dilute nitric acid to the third portion of the solution followed by about 1 cm ³ of barium nitrate solution.	[2]
(f)	Add about 1 cm ³ of dilute nitric acid to the fourth portion of the solution followed by about 1 cm ³ of silver nitrate solution.	

(g)	What does test (f) tell you about solid U?	
		[1]
(h)	Name the gas given off in test (c).	
		[1]
(i)	What conclusions can you draw about solid U ?	
		[3]
		[-]
	[Total:	18]

BLANK PAGE

BLANK PAGE

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

anion	test	test result
carbonate (CO ₃ ^{2–})	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I [_]) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO₃⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ^{2–}) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al ³⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ +)	ammonia produced on warming	-
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results
ammonia (NH ₃)	turns damp red litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl_2)	bleaches damp litmus paper
hydrogen (H ₂)	'pops' with a lighted splint
oxygen (O ₂)	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.