## **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**International General Certificate of Secondary Education** 

## MARK SCHEME for the October/November 2012 series

## 0620 CHEMISTRY

0620/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2012	0620	31

A - A-V - Affiliation on the attenual attaining	
1 (a) diffusion or fractional distillation;	

	` '		
	(b) frac	ctional distillation;	
	(c) sim	ple distillation;	
	(d) cry	stallisation;	
	(e) filtr	ation;	
	(f) chr	omatography;	
			[Total: 6]
2	(a) (i)	become darker;	[1]
	(ii)	increase;	[1]
	(iii)	black / dark grey;	[1]
		not: brown solid;	[1]
	(b) (i)	same Z / same number of protons; accept: atoms of the same element	[1]
		different number of neutrons / different nucleon number / different mass number;	[1]
	(ii)	53 protons and 53 electrons;	[1]
		78 neutrons;	[1]
	(iii)	xenon;	[1]
		F <sub>3</sub> / F <sub>3</sub> Br; F <sub>5</sub> / F <sub>5</sub> Br;	[1] [1]
			[Total: 11]

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2012	0620	31
3 (a) (i)	any three from:		

(a) (i)	any three from: particles have more energy; move faster; collide more frequently; more successful collisions; accept: atoms or molecules for particles not: electrons not: vibrate more	[3]
(ii)	reaction faster with temperature increase; enzymes denatured / destroyed; <b>not:</b> killed	[1] [1]
(b) (i)	bigger initial gradient; same final volume of nitrogen;	[1] [1]
(ii)	decrease / slows down;	[1]
(iii)	<pre>concentration of organic compound decreases; compound used up = [1] or: fewer particles; collision rate decreases;</pre>	[2]
(c) (i)	carbon monoxide-incomplete combustion; carbon - containing fuel / fossil fuel / petrol;	[1] [1]
	oxides of nitrogen - oxygen and nitrogen react; at high temperature / in engine; <b>not</b> : in exhaust	[1] [1]
(ii)	carbon monoxide to carbon dioxide; oxides of nitrogen to nitrogen; correct balanced equation;	[1] [1] [1]
		[Total: 17]

	Pa	Page 4		Mark Scheme		Paper	
				IGCSE – October/November 2012	0620	31	
4	(a)	•	nt covalent; polymer made from monomers;				
	(b)	(i)	high hard brittle insol	•		[3]	
		(ii)		on / diamond / silicon / boron; graphite		[1]	
	(c)	(i)	sodi	um hydroxide / any named alkali / reactive metal;		[1]	
		(ii)		ed acid; onium oxide;		[1] [1]	
						[Total: 8]	
5	(a)	(i)	influe or: turns	of reaction; enced by light / only happens in light; s light into chemical energy = [2] ept: light is catalyst = [1]		[1] [1]	
		(ii)	they appr or: phot corredioxi anyt effect or: chlor mak	are reduced to silver / 2AgCl → 2Ag + Cl₂; ropriate importance given; rosynthesis; rect comment about chemistry carbon dioxide to carbide to oxygen; hing sensible e.g. its role in the food chain or decreated or oxygen for respiration; rination; ring chloroalkanes; ropriate importance given;	•	[1] [1] [1]	
	(b)	(i)		sure would move position of equilibrium to right / incease pressure favours side with less (gas) molecules		[1] [1]	
		(ii)		ease temperature favours endothermic reaction; ess products/reduce yield;		[1] [1]	
		(iii)	keep	os rate high / increase rate at lower temperatures;		[1]	

	Page 5			Mark Scheme	Syllabus	Paper
				IGCSE – October/November 2012	0620	31
	(c)	eac 4 e 2 nt	[1] [1] [1]			
				[Total: 13]		
6	(a)	(i)	ol;	[1] [1] [1]		
		(ii)	poly	ester;		[1]
		. ,	allo	<b>w:</b> named polyester amide;		[1]
				w: nylon		[1]
	(b)			ect amide linkage; amide linkage correctly orientated		[1]
		– N		[1]		
		not	e: mo	onomers are amino acids not diamines or dicarboxy	lic acid	
	(c)	bro		[1]		
		uns satı	[1] [1]			
		or:				
		or:	from	ic potassium manganate(VII) purple/pink to colourless; <b>not:</b> clear s purple;		
						[Total: 10]
7	(a)	(i)		ing point is below 25°C; ng point above 25°C;		[1] [1]
			acce	ept: argument based on actual values e: 25°C is between mp and bp = [2]		[1]
		(ii)		ntium loses 2e; ır gains 2e;		[1] [1]
		(iii)		rogen chloride / hydrochloric acid;		[1]
		···· <i>)</i>	-	ept: sulfurous acid or sulfur dioxide		[.]
		(iv)		en strontium chloride has ions/ionic compound;		[1] [1]
				ur chloride has no ions / only molecules / molecular	/ covalent;	[1]

(b) (i)	strontium carbonate does not dissolve / no effervescence; note: not just reaction is complete	[1]	
(ii)	to remove excess/unreacted / undissolved strontium carbonate;	[1]	
(iii)	water of crystallisation needed / $6H_2O$ in crystals / would get anhydrous salt / would not get hydrated salt / crystals dehydrate; <b>not</b> : just to obtain crystals	[1]	
(c) number of moles of HC $l$ used = 0.05 × 2 = 0.1 number of moles of SrC $l_2$ .6 H $_2$ O which could be formed. = 0.05 mass of one mole of SrC $l_2$ .6H $_2$ O is 267 g theoretical yield of SrC $l_2$ .6H $_2$ O = 0.05 × 267 = 13.35 g percentage yield = 6.4/13.35 × 100 = 47.9% accept: 48% allow: ecf			

Mark Scheme
IGCSE – October/November 2012

Page 6

Syllabus 0620

[Total: 15]

Paper 31