UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

General Certificate of Education O Level

MARK SCHEME for the June 2004 question papers

	5054 PHYSICS
5054/01	Paper 1 (Multiple Choice), maximum mark 40
5054/02	Paper 2 (Theory), maximum mark 75
5054/03	Paper 3 (Practical Test), maximum mark 30
5054/04	Paper 4 (Alternative to Practical), maximum mark 30

These mark schemes are published as an aid to teachers and students, to indicate the requirements of the examination. They show the basis on which Examiners were initially instructed to award marks. They do not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2004 question papers for most IGCSE and GCE Advanced Level syllabuses.



MARK SCHEME

MAXIMUM MARK: 40

SYLLABUS/COMPONENT: 5054/01

PHYSICS
Paper 1 (Multiple Choice)

Page 1	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	1

Question Number	Key	Question Number	Key
1	Α	21	Α
2	Α	22	D
3	С	23	С
4	D	24	Α
5	С	25	С
6	D	26	В
7	С	27	В
8	Α	28	D
9	D	29	В
10	Α	30	Α
11	С	31	В
12	В	32	D
13	Α	33	D
14	Α	34	В
15	D	35	С
16	В	36	Α
17	В	37	Α
18	В	38	D
19	D	39	Α
20	С	40	В

TOTAL 40

MARK SCHEME

MAXIMUM MARK: 75

SYLLABUS/COMPONENT: 5054/02

PHYSICS Paper 2 (Theory)



Page 1	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	2

Section A

1	(a)	(i)	weight / gravity / gravitational (force)		
		(ii)	air / wind resistance or drag or friction / upthrust		B1
	(b)	(i) (ii)	9.8 or 10 m/s ² or N/kg air resistance increases (not if parachute open) less resultant force or sensible statement about upwards force e.g. resistance opposes gravity or decreases acc. slope of line decreases	C1 C1 C1	B1 B2
		(iii)	air resistance = weight / no resultant / net / overall force / down	ıwards	
		, ,	force balances upwards force	Total	B1 [6]
2	(a)	(i)	radiation		B1
	/b\	(ii)	no molecules or medium (to vibrate, conduct, convect) / vacuum hot air rises		B1 B1
	(b)		(hot) air expands / density decreases		B1
	(c)		fiberglass or air is a bad conductor/ insulator / lags / reduces hea	at flow	D4
			fiberglass traps air or prevents convection (ignore radiation statements)		B1 B1
				Total	[6]
3	(a)		rise in temperature / hot / heated		B1
	()		road / bridge / rail / metal expands or gap reduces		B1
	(b)		no buckling / deformation / breaking / cracking / twisting / tilting any other problem + solution e.g. concrete cracks – leave a gap, telephone wires sag – put them high / tight hot water cracks glass – use thin glass / car engines seize up – cool them water freezes in pipes – lag them or use antifreeze / tyres burst – let air out pipes bend – use flexible joints / dashboard deforms – car in shade		B1
			wrong readings on measuring cylinder – use correct temp.		B1
				Total	[4]
4	(a)		distance traveled per unit time or in one second / distance ÷ time	•	5.4
	(b)		or rate of change of distances = d/t in any algebraic or numerical form		B1 C1
	` '		any doubling of distance or final time		C1
	(c)		0.48 s (allow 0.24s 2/3 accept 0.5s) 60/0.48 (5)		A1 C1
	(-)		123.75 accept 120, 123, 124 (ecf (b))		A1
				Total	[6]
5	(a)	(i)	magnetic (field) of current / coil / recording head		
		(ii)	or head is magnetized / an electromagnet magnetism / magnetic field or current or poles on head reverses	1	B1
			changes direction (accept "due to alternating current")	,	В1
	(h)	(iii)	each direction / one cycle longer (on tape)		B1 B1
	(b)	(i) (ii)	need to keep record / tape stored or played iron, steel etc		B1
		- *		Total	[5]

	Page 2		Mark Scheme	Syllabus	Paper	
			PHYSICS – JUNE 2004	5054	2	
6	6 (a) (voltage past maximum or 3V / off scale / outside reading less accurate or sensitive / not far up scadeflection	•	r	B1
	(b)	(i)	V = I R in any algebraic format 4/12 0.33 A (accept 1/3 A)			B1 C1 A1
		(ii)	(i) * 30 or (i) * 18 + 4 or 30*4/12 9.9 - 10 V (e.c.f (i), e.g. if (i) = 0.3, 0.3*30 = 9V or 0.3*18+4 only 1 unit error in this question	4 = 9.4 V)		C1 A1
			·		Total	[7]
7	(a)	(i)	filament is hot / heated (by current from 6V supplemission	y) / thermior	ic	B1
		(ii)	anode is positive / anode attracts electrons / elec	trons attract	ed to +	B1
		(iii)	(electric) field from anode to cathode otherwise electrons stopped / deflected / slowed	down /		
			collide (with air atoms) (accept no opposition to movement, to reach screen, to avoid air re	sistance)		B1
	(b)		up and down vertical or side to side movement electrons deflected by electric field or attracted to	(not on both	,	В1
			or plates are charged (e.g. plates are +ve and -v	•	tu by –	В1
					Total	[5]
8	(a)		radon (gas) cancer / mutation / cell damage or death			B1
	(b)		radiation sickness or adds to readings			В1
	(c) (d) (e)		(accept count with no source) (outer) space / stars / Sun number of protons and neutrons (not no. nucleons) 84 216 (values reversed B1)	ot sunlight)	Total	B1 B1 B2 [6]

Mark Scheme	Syllabus	Paper
PHYSICS – JUNE 2004	5054	2

SECTION B

9	(a)	(i)	Any three other parts of spectrum radio, microwaves, u.v., X, γ (-1 any wrong if>3 ignore t.v.)	М3
			correct order for all including visible (accept colours) and I.R.	A1
		(ii)	reflection of infra-red or radiation (from shiny material)	B1
		. ,	more energy hits food or reflection towards food	
			cooks food faster	
			avoids wasting heat / energy or more efficient	
			avoids heating outer case or burning hand ANY 2	B2
	(b)		connected to (outer metal) case	В1
	• •		if live touches case or case becomes live	B1
			allows current / charge to earth / ground	B1
			blows fuse (and disconnects circuit)	
			or no current through person or no electrocution / electric shock	B1
	(c)	(i)	P = V I in any algebraic form	В1
	. ,	(ii)	230 * 8.3	C1
			1900 W (accept 1910 W but not power 1/4)	A1
		(iii)	current decreases (halves) or power 1/4	B1
			Total	[15]
10	(a)		mass of bar (measured)	M1
			using (top-pan) balance / spring balance / scales etc.	A1
			length, breadth and height measured	
			or volume water + bar measured or displacement can (full) with water volume = length x breadth x height	M1
			or subtract volume water alone or collect water displaced	A1
			using ruler / calipers / micrometer or measuring cylinder	A1
			density = mass / volume	B1
	(b)) (i)	melts / changes state / becomes liquid	B1
	(-)	(ii)	(initial) increase in vibration / K.E. of molecules (to 600s)	B1
		()	then later / after 600s or on melting	
			bonds broken (accept molecules break free / overcome attraction / not fixed in place)	B1
		(iii)	$E = mc (\Delta)T$ algebraic form seen	C1
			645 – 655 (°C) seen)	C1
			17 160 J (allow 1700, 17200, 20000)	A1
		(iv)	30*400 or 12 000 (J) seen)	C1
			E = mL any algebraic form seen or 12 000/0.3	C1
			40 000 J/kg	A1
			Total	[15]

Page 4	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	2

11	(a)	(i) (ii) (iii)	P.E. decreases (A to B or C to D or downhill or initially) K.E. gained (P.E. \rightarrow K.E2) K.E. to P.E. change must be clear and from B to C or uphill mgh algebraic form seen 500*10*30 150 000 J conservation of energy cited or clear that loss of P.E. has become K.E. 500*10*20 or 500*10*10 or 50 000 seen 100 000 J (allow g=9.8)	B1 B1 C1 C1 A1 C1 A1
	(b)	(i) (ii)	velocity involves direction or is a vector (speed does not) direction (of carriage) changes / carriage turns (accept on diagram) force towards centre (of curve) / inwards (accept centripetal)	B1 B1
	(c)		F = ma in any algebraic form or 3000 = 500a 3000/500 6(.0) m/s ² Total	C1 C1 A1 [15]
			Total for paper :	[75]

MARK SCHEME

MAXIMUM MARK: 30

SYLLABUS/COMPONENT: 5054/03

PHYSICS
Paper 3 (Practical Test)



1.	(a), (b) & (c)	Repeat measurements taken for either t_1 or t_2 .	B1
		Correct T_1 in the range 1.40 s to 1.60 s to 0.01 s	B1
		Correct T_2 within ± 0.1 s of T_1	B1
	(d)	Comment on Either reaction time – however expressed Or range of values	B1
	(e) Or	Sensible conclusion based on their results e.g. Time for one oscillation is independent of the mass. (if periods are the same within the limits of uncertainty) Time for one oscillation increases / decreases with increase in mass. (Allow direct or inverse proportion) (provided their results show this)	B1
		Tota	ıl [5]
2.	(a)	Power supply, ammeter and switch in series with gap between A and B, voltmeter in parallel with power supply.	
2.	(a) (b), (c) & (d)	Power supply, ammeter and switch in series with gap between A and B, voltmeter in parallel with power supply.	
2.	. ,	Power supply, ammeter and switch in series with gap between A and B, voltmeter in parallel with power supply. I values in region of 0.3 A and 0.45 A with unit seen at least once and at least one current to 0.01 A.	B1
2.	. ,	Power supply, ammeter and switch in series with gap between A and B, voltmeter in parallel with power supply. I values in region of 0.3 A and 0.45 A with unit seen at least once and at least one current to 0.01 A. (Allow Centre variation) Both V values in the region of 4.5 V with unit seen at least once and at least one voltage to 0.1 V.	B1
2.	. ,	Power supply, ammeter and switch in series with gap between A and B, voltmeter in parallel with power supply. I values in region of 0.3 A and 0.45 A with unit seen at least once and at least one current to 0.01 A. (Allow Centre variation) Both V values in the region of 4.5 V with unit seen at least once and at least one voltage to 0.1 V. (Allow Centre variation) R values in the region of 15 Ω and 10 Ω with unit seen at	B1 B1 B1 B1

Mark Scheme PHYSICS – JUNE 2004

Page 1

Syllabus 5054 Paper 3

P	age 2			Mark Scheme	Syllabus	Paper
			P	HYSICS – JUNE 2004	5054	3
3.	(a) an	d (b)	Sensible	temperatures with unit seen at least	once.	B1
			At least o	one reading attempted to better than 1 $^{\circ}$	С	B1
			and corr	erically to (1.0 to 3.0) x temperature ect calculation of $V_{\rm I}$ with unit seen a crically equal to $V_{\rm I}$.	-	B1
	(c) an	d (d)		values for all the thermal energy ch at least once.	anges with	M1
	(e)			gained greater than energy lost as co energy from beaker / surroundings	ld water gain	ns A1
					To	otal [5]
4. <u>In</u>	<u>itial rea</u>	dings.				
	(b)		<i>x</i> 0.60 <u>+</u>	0.05 m with unit.		B1
	(c)		_	0.05 m with unit. e missing unit once only)		B1
Т	<u>able</u>		•	recorded to 0.001 m or better.		B1
	abic					
	(d)		Table w	ith units for d , D and $1/D$.		B1
			At least	one reading with D greater than or e	equal to 1.00	m. B1
			At least	one reading with D less than or equa	al to 0.70 m	B1
			Correct	calculation of $(d/D)^2$ and $1/D$ to at 1	east 2 s.f.	B1
			Five goo	od values judged according to the tal	ole below.	B1
		D		Range of $(d/D)^2$	1 /	\overline{D}
				<i>U</i> \ /		

D	Range of $(d/D)^2$	1/D
0.65	0.06 - 0.10	1.54
0.70	0.12 - 0.16	1.43
0.75	0.18 - 0.22	1.33
0.80	0.23 - 0.27	1.25
0.85	0.27 - 0.31	1.18
0.90	0.31 - 0.35	1.11
0.95	0.35 - 0.39	1.05
1.00	0.38 - 0.42	1.00

Page 3	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	3

Graph.

(e) Axes labelled with unit and correct orientation. B1 Suitable scale y axis 1 cm = 0.02 / 0.025x axis 1 cm = 0.1 or 0.05 m⁻¹ M1 Two points plotted correctly – check the two points furthest from the line. **A**1 Best fit fine line and finely plotted points. Β1 **Calculations.** (f) and (g) Large triangle. B1 Correct calculation of *S* and *f* (ignore sign) Β1 Value of f in range 0.130 m to 0.170 m with unit. B1

MARK SCHEME

MAXIMUM MARK: 30

SYLLABUS/COMPONENT: 5054/04

PHYSICS (Alternative to Practical)



Page 1	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	4
Question 1			
One X an	rays from X and Y (clear <u>intention</u> to touch hole edges d one Y ray "touch" an edge of the hole and meet scre X and one Y are neat lines (rule and sharp "pencil") allo	én	M1 C1
"refraction" or "diffraction" at hole One correct X and the corresponding Y labeled on screen			B1
Arrows or	rays; no broken lines penalty -1 (max).		B1
(b) XY in ran	ge 54 to 56 mm (unit required), accept in cm		B1
		To	tal [5]
Question 2			
Accept ar comprom		rcuit is not	
Penalise	-1 (max) :- short circuit (e.g. line behind component, ur	nless signs (of use

(b) Correct polarities, +ve signs for correct terminals of cell and ammeter (re diode).

of rubber) or any compromised circuit function.

(c) No current / I = 0, (do not accept "nothing"), accept very small "reverse" current / lamp does not light.

(d) One from: limit current / prevent overheating / current indicator / provides resistance

Total [6]

В1

В1

B1

В1

В1

B1

Question 3

(a) Any method <u>based</u> on rule reading at 25°C – rule reading at top of thermometer bulb.
 NB / required. Mark text or diagram or <u>Fig 3.1</u>
 Rule as close as possible to thermometer (on diagram < 1 cm) / uses fiducial aid

With the eye/line of sight perpendicular to the rule/end of mercury thread B1

(b) (i) $I_0 = 5.6 - 5.8$ (cm), $I_{100} = 22.6 - 22.8$ (cm) ignore unit (ii) $\Delta I / 100$, clear, correct arithmetic ecf, 2 or 3 dcp, ignore unit, accept any correct $\Delta I / \Delta \theta$ from graph.

(iii) linearly, or $(I - I_0) \propto \theta$ accept/line has a constant/uniform m, note that... "directly proportional" automatically looses the mark.

Total [6]

Page 2	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	4

Question 4

(a) (i) V initial = a volume between 40cm ³ and 60cm ³ : (allow use of beaker) must be able to displace 40cm ³ / prevents overflowing /	B1		
exceeding cm ³ limit	B1		
(ii) $\{V_{\text{max}} - V_{\text{initial}}\}$ / change in volume is found / change in volume obtained is = V_{metal} / any related answer that has an association of measurement of volume.	B1		
 (iii) Any good point e.g. tap cylinder to release air / how avoiding parallax / water at 20°C / careful pouring / avoid splashing / use set square / repeat average / reading the position of the bottom of the meniscus. (b) Scale calibration of cylinder is correct at 20°C / liquid needs to be at 20°C 	B1 B1		
(c) Water (on the metal would be) included in the (repeat) volume of the metal; or something that means the same, not just erroneous.	В1		
Total [6]			
Question 5			
(a) Axes correct, scale that cannot be x2 / is not "awkward" and with units Correct plotting, nearest ½ small square, check first point and obvious			
plot errors. Line judgement re plots (line does not go through all correctly plotted points,	B1		
so accept smooth line through 5 points i.e., one point not on the line) Neat smooth thin line			
(b) Mark cands diagram or Fig 5.1: (i) Object displace downwards OR screen displaced downwards	B1		
Any ray from the top of object through the lens to meet screen.	В1		
Be generous re art and accuracy of position, (ii) put centres in line	B1		
Tota	I [7]		
Paper Total	30		