MARK SCHEME for the October/November 2014 series

0625 PHYSICS

0625/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.



Page 2	Mark Scheme Cambridge IGCSE – October/November 2014	Syllabus 0625	Paper 31	
		0025	31	
	NOTES ABOUT MARK SCHEME SYMBOLS AND OTHER MATTERS			
B marks	B marks are independent marks, which do not depend on other marks. For a B mark to be scored, the point to which it refers must be seen specifically in the candidate's answer.			
M marks	M marks are method marks upon which accuracy marks (A marks) M mark to be scored, the point to which it refers must be seen in a If a candidate fails to score a particular M mark, then none of the c can be scored.	a candidate's	s answer.	
C marks	C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, provided subsequent working gives evidence that they must have known it . For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.			
A marks	A marks are accuracy or answer marks which either depend on an one of the ways which allow a C mark to be scored. A marks are c final answers to numerical questions. If a final numerical answer, e correct, with the correct unit and an acceptable number of significa- marks for that question are normally awarded. It is very occasiona a correct answer by an entirely wrong approach. In these rare circl award the A marks, but award C marks on their merits. An A mark a dependent mark.	ommonly av eligible for A ant figures, a lly possible t umstances,	varded for marks, is Il the o arrive at do not	
Brackets ()) Brackets around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.			
<u>Underlining</u>	<u>q</u> Underlining indicates that this must be seen in the answer offered, or something very similar.			
OR / or	This indicates alternative answers, any one of which is satisfactory	/ for scoring	the marks.	
e.e.o.o.	This means "each error or omission".			
o.w.t.t.e.	This means "or words to that effect".			
Ignore	This indicates that something which is not correct or irrelevant is to does not cause a right plus wrong penalty.	be disrega	rded and	

- Spelling Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, do not allow ambiguities, e.g. spelling which suggests confusion between reflection / refraction / diffraction or thermistor / transformer.
- Not / NOT This indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate, i.e. right plus wrong penalty applies.

Page 3	Mark Scheme	Syllabus	Paper	
	Cambridge IGCSE – October/November 2014	0625	31	
ecf	meaning "error carried forward" is mainly applicable to numerical questions, but may in particular circumstances be applied in non-numerical questions. This indicates that if a candidate has made an earlier mistake and has carried an incorrect value forward to subsequent stages of working, marks indicated by ecf may be awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate from being penalised more than once for a particular mistake, but only applies to marks annotated ecf.			
Sig. figs.	Answers are normally acceptable to any number of significant figure exceptions to this general rule will be specified in the mark scheme the second or third significant figure will be penalised.			
Arithmetic errors Deduct one mark if the only error in arriving at a final answer is clearly an arithmetic one. Regard a power-of-ten error as an arithmetic error.				
Transcription errors Deduct one mark if the only error in arriving at a final answer is because previously calculated data has clearly been misread but used correctly.				

- Fractions Allow fractions only where specified in the mark scheme.
- Units Deduct one mark for an incorrect or missing unit, but only if the answer would otherwise have gained all the marks available for that answer. Maximum one unit penalty per question.

Page 4		Mark Scheme		Paper
		Cambridge IGCSE – October/November 2014	0625	31
1	B C	increasing speed constant speed stationary ote: one mark lost for e.e.o.o.		B2
	(b) D E F No		B2	
	(c) (i)	$(a =) \Delta v/t \text{ OR } (v-u)/t \text{ OR } 10.5/1.5$ = 7.0 m/s ²		C1 A1
	(ii)	(<i>a</i> =) 0 (m/s ²)		B1
	(iii)	upward and downward forces equal OR no resultant force OR forces equal and opposite OR forces balanced OR weight (of body) = tension (in rope)		B1 [Total: 8]
2	(a) (i)	(increase in g.p.e. = mgh OR $65 \times 10 \times 8 =$) 5200 J		B1
	(ii)	EITHER k.e. gained = g.p.e. lost $\frac{1}{2} mv^2$ = 5200 in any form v^2 = 5200/(0.5 × 65) OR 160 v = 12.6 m/s e.c.f. (a)(i) OR		C1 C1 C1 A1
		$v^{2} = u^{2} + 2as/v^{2} = 2gh$ $v^{2} = 2 \times 10 \times 8$ $v^{2} = 160$ v = 12.6 m/s e.c.f. (a)(i)		(C1) (C1) (C1) (A1)
	• • •	eed is the same THER		B1
	los	is in g.p.e. is the same a. gained is the same		B1 B1
	ac	celeration is the same stance fallen is the same		(B1) (B1)
				[Total: 8]

P	age	5	Mark Scheme	Syllabus	Paper
	ago	•	Cambridge IGCSE – October/November 2014	0625	31
3	(a)	(i)	force/pressure greater on outside surface owtte		B1
		(ii)	p = F/A in any form OR ($F =$) pA		C1
			= $(1.0 \times 10^5 - 6000) \times 0.12$ 11280 N to at least 2 sig. figs.		C1 A1
			1 1200 N to at least 2 sig. ligs.		AI
	(b)	(i)	pressure of oil = pressure of water		B1
		(ii)	1. (<i>p</i> =) <i>hρg</i> (= 0.25 × 1000 × 10 =) 2500 Pa		C1 A1
			2. $h\rho g = 2500$ ($\rho = 2500/(0.32 \times 10) =$) 781 kg/m ³ to at least 2 sig. figs.		C1 A1
					[Total: 9]
4	(a)		ss of block m		B1
			al temperature θ_1 and final temperature θ_2 e of heating <i>t</i>		B1 B1
			age/p.d. V AND current I		B1
	(b)		•) $VIt \div [m (\theta_2 - \theta_1)]$ $Pt \div [m (\theta_2 - \theta_1)]$ OR $E \div [m (\theta_2 - \theta_1)]$ as appropriate to symbols defined.	fined in (a)	
		nur	nerator correct		B1
		der	nominator correct		B1
	(c)	(mo	ore) thermal energy/heat lost (to surroundings) so temperature rise i	<u>s less</u>	
		OR	more thermal energy/heat input required for same temperature rise	<u>e</u>	B1
					[Total: 7]
5	(2)	(i)	longitudinal: oscillations/vibration of particles/molecules in direction	of travel	
U	(u)	(')	(of wave)		B1
			transverse: oscillation/vibrations of particles/molecules perpendicul direction of travel (of wave)	lar to	B1
		(ii)	1. e.g. sound wave / compression wave on a spring		B1
			2. e.g. any named electromagnetic wave / ripples / water wave / water the stretched rope	wave on a	B1
	(b)		e of $v = f\lambda$ in any form OR $(\lambda =) v/f$ OR 7200/30 OR 7.2/30		C1
		240)m / 0.24 km		A1
	(c)	no	no sound heard/quieter sound		B1
	. ,	me	dium/air required to transmit sound sound does not travel through a vacuum		B1
			Sound does not traver through a vacuum		
					[Total: 8]

6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0625	31
(i)	 one normal to mirror drawn angle of incidence, labelled X 		B1 B1
(ii)	 both reflected rays drawn construction lines to locate image, marked I 		B1 B1
(i)	dot marked C in correct position		B1
(ii)	two circular arcs each joining correct points on barrier spacing of arcs same as spacing of incident waves		B1 B1
			[Total: 7]
(i)	diagram showing: molecules widely spaced molecules randomly positioned		B1 B1
(ii)	(attractive) forces (much) smaller between gas molecules gas molecules (much) farther apart		B1 B1
(i)	pV = constant OR $p_1V_1 = p_2V_2$ OR $(V_2 =) p_1V_1/p_2$ OR $(V_2 =) 2.75 \times 10^6 \times 6 \times 10^{-3}/1.1 \times 10^5$ = 0.15 m ³ (no. of balloons = $(0.15 - 6 \times 10^{-3})/3 \times 10^{-3}$ =) 48		C1 C1 A1
(ii)	pressure of air in balloon increases molecules move faster OR hit balloon surface harder/more often OR larger force rips/breaks rubber OR balloon expands		B1 B1
			[Total: 9]
(i)	rectifier/diode		B1
(ii)	frequency (of A.C. supply)		B1
(i)	(<i>P</i> =) <i>IV</i> OR 0.5 × 5.3 OR 500 × 5.3 2.6 W OR 2600 mW		C1 A1
(ii)	(<i>E</i> =) <i>Pt</i> OR <i>IVt</i> OR $2.65 \times 1.5 \times 3600$ OR $0.5 \times 5.3 \times 1.5 \times 3600$ 14000 J	0	C1 A1
ene	ergy only underlined		B1
			[Total: 7]
	 (ii) (i) 	 Cambridge IGCSE – October/November 2014 (i) 1. one normal to mirror drawn 2. angle of incidence, labelled X (ii) 1. both reflected rays drawn 2. construction lines to locate image, marked I (i) dot marked C in correct position (ii) two circular arcs each joining correct points on barrier spacing of arcs same as spacing of incident waves (i) diagram showing: molecules widely spaced molecules randomly positioned (ii) (attractive) forces (much) smaller between gas molecules gas molecules (much) farther apart (i) <i>pV</i> = constant OR <i>p</i>₁<i>V</i>₁ = <i>p</i>₂<i>V</i>₂ OR (<i>V</i>₂ =) <i>p</i>₁<i>V</i>₁<i>/p</i>₂ OR (<i>V</i>₂ =) 2.75 × 10⁶ × 6 × 10⁻³/1.1 × 10⁵ = 0.15 m³ (no. of balloons = (0.15 - 6 × 10⁻³)/3 × 10⁻³ =) 48 (ii) pressure of air in balloon increases molecules move faster OR hit balloon surface harder/more often OR larger force rips/breaks rubber OR balloon expands (i) rectifier/diode (ii) frequency (of A.C. supply) (ii) (<i>P</i> =) <i>IV</i> OR 0.5 × 5.3 OR 500 × 5.3 2.6W OR 2600 mW (ii) (<i>E</i> =) <i>Pt</i> OR <i>IVt</i> OR 2.65 × 1.5 × 3600 OR 0.5 × 5.3 × 1.5 × 3600 	Cambridge IGCSE - October/November 2014 0625 (i) 1. one normal to mirror drawn 2. angle of incidence, labelled X (ii) 1. both reflected rays drawn 2. construction lines to locate image, marked I (i) dot marked C in correct position (ii) two circular arcs each joining correct points on barrier spacing of arcs same as spacing of incident waves (i) diagram showing: molecules widely spaced molecules randomly positioned (ii) (attractive) forces (much) smaller between gas molecules gas molecules (much) farther apart (i) $pV = \text{constant OR } p_1V_1 = p_2V_2 \text{ OR } (V_2 =) p_1V_1/p_2$ OR $(V_2 =) 2.75 \times 10^6 \times 6 \times 10^{-3}/1.1 \times 10^5$ = 0.15 m ³ (no. of balloons = $(0.15 - 6 \times 10^{-3})/3 \times 10^{-3} =) 48$ (ii) pressure of air in balloon increases molecules move faster OR hit balloon surface harder/more often OR larger force rips/breaks rubber OR balloon expands (i) rectifier/diode (ii) frequency (of A.C. supply) (i) $(P =) IV$ OR 0.5×5.3 OR 500×5.3 2.6W OR 2600 mW (ii) $(E =) Pt$ OR IVt OR $2.65 \times 1.5 \times 3600$ OR $0.5 \times 5.3 \times 1.5 \times 3600$

Pa	age T	7	Mark Scheme	Syllabus	Paper
			Cambridge IGCSE – October/November 2014	0625	31
9	(a)		ekground (radiation) OR a specific source of background radiation ending materials/radon gas/cosmic rays	e.g. rocks/	B1
	(b)	low slig ver	r three from: count rate due to background radiation only htly less reading due to random nature of radioactivity y high reading due to α-particles OR emission from source Iden increase of count rate at limit of range of α-particles		В3
	(c)	(i)	downward <u>curve</u>		B1
		(ii)	(count rate) decreases/background only deviation starts at start of plates		B1 B1
					[Total: 7]
10	(a)	 (lamps) stay on/have same brightness as before/nothing happens (lamps) still connected to supply/have same voltage as before/are connected in parallel 			B1
	()			nected in	B1
	(b)	(i)	line 1: on line 2: off line 3: off line 4: on deduct one mark for e.e.o.e.		B2
		(ii)	when either switch is operated, the state of the lamp changes.		B1
					[Total: 5]
11	(a)	(i)	electromagnetic induction		B1
	(b)	(i)	pointer deflects pointer returns to zero		B1 B1
		(ii)			B1
			pointer deflects in opposite direction <u>and</u> returns to zero OR deflects for shorter time		B1
					[Total: 5]