## MARK SCHEME for the May/June 2015 series

## **5070 CHEMISTRY**

5070/41

Paper 4 (Alternative to Practical), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.



Ρ	age 2	Mark Scheme	Syllabus	Paper
		Cambridge O Level – May/June 2015	5070	41
1	(a) (i)	Silver/grey solid (1)		[1]
	(ii)	$2Zn + O_2 = 2ZnO(1)$		[1]
	(iii)	Nitric acid/HNO <sub>3</sub> (1)		[1]
	(b) (i)	Toxic/poisonous gas evolved (1)		[1]
	(ii)	3.78/189 = 0.02 (1)		[1]
	(iii)	$0.02 \times 24000 \times 2 = 960 \text{ cm}^3$ (1) $0.02 \times 24000 \times 0.5 = 240 \text{ cm}^3$ (1)		[2]
				[Total: 7]
2	<b>(a)</b> 88 n	8 – 45 = 43 (1) = 3 (1)		
	bı	utanoic acid/butyric acid (1)		[3]
	<b>(b)</b> hy po	rdrogen (1) ops in flame / burning splint pops / lighted splint pops (1)		[2]
	(c) (i)	esters (1)		[1]
	(ii)	$CH_3COOC_2H_5$ / $CH_3CO_2C_2H_5$ (1)		[1]
	(iii)	ethanol (1) ethanoic acid (1)		[2]
	(iv)	$C_2H_5COOCH_3(1)/methyl propanoate (1)$		
		<b>OR</b> HCOOC <sub>3</sub> H <sub>7</sub> (1)/propyl methanoate (1)		[2]
				[Total: 11]
3	(d)			[Total: 1]
4	(b)			[Total: 1]
5	(b)			[Total: 1]
6	(b)			[Total: 1]
7	(b)			[Total: 1]

Ρ	age 3	3	Mark Scheme		Syllabus	Paper
			Cambridge O Level – May/Ju	ne 2015	5070	41
8	(a)	Ga	s no longer turns litmus blue (1)			[1]
	(b)	Pin	k to colourless (1)			[1]
	(c)	29 <u>8</u> 20	.5 28.8 39.9 1 mark for each correc .9 <u>7.9 19.5</u> to the benefit of the ca .6 <u>20.9 20.4</u>	t row <u>or</u> column ndidate (3)		
		Ме	an value 20.5 cm³ (1)			[4]
	(d)	0.0	0205 moles (1)			[1]
	()					[.]
	(e)	(i)	0.00205 (1)			[1]
		(ii)	0.0205 (1)			[1]
	(f)	0.1	(1)			[1]
	(g)	0.0	795 (1)			[1]
	(h)	(i)	1.352 g (1)			[1]
	()	(ii)	54 06 g (1)			[1]
		(11)	34.00 g (1)			[']
	(i)	On	e mole of $(NH_4)_2SO_4$ produces $34g/2$ moles	of ammonia (1)		
		Co	ncentration = $54.06/34 = 1.59 \text{ mol}/\text{dm}^3$ (1)			[2]
						[Total: 15]
٥	(2)	col	ourloss solution (1)			
3	(a)	CON				
	(b)	(i)	white ppt (1)			
		(ii)	soluble in excess (1)			
	(c)	(i)	white ppt (1)			
		(ii)	insoluble in excess (1)			
	(d)	M1	(aq) NaOH/sodium hydroxide/ (1)			

M2 Al/aluminium (foil)/Devarda's alloy (1) M3 warm/heat/boil (1) M4 ammonia / NH<sub>3</sub> OR gas turns litmus blue (1)

Page 4		Mark Scheme	Syllabus	Paper
		Cambridge O Level – May/June 2015	5070	41
		<b>ALLOW</b> Brown ring test: conc. (1) sulfuric acid/H <sub>2</sub> SO <sub>4</sub> (1) iron(II) sulfate/FeSO <sub>4</sub>	1 (1) brown	ring (1) <b>[Total: 9]</b>
10	(a)	10, 36, 54, 68 All correct for two marks; three correct for one mark		[2]
	(b)	Temperature at which solid appears is below room temperature (1) Cooling the tube by some method e.g. ice (1)		[2]
	(c)	all points plotted correctly (1) two smooth curves through the points (1 mark for each)		[3]
	(d)	(i) $NH_4Cl - 2.8$ (1)		[1]
		(ii) KNO <sub>3</sub> – 1.7 (1)		[1]
	(e)	(i) 23 (1)		[1]
		(ii) 3.4 g/10g = 34 g/100 g water (1)		[1]
	(f)	$NH_4Cl$ – solution + undissolved solid (1) KNO <sub>3</sub> – solution (no solid) (1)		[2]
				[Total: 13]