# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education

Advanced Subsidiary Level and Advanced Level

#### **CHEMISTRY**

Paper 2 Structured Questions AS Core



9701/02

May/June 2006

1 hour 15 minutes

Candidates answer on the Question Paper. Additional Materials: Data Booklet

Candidate Name							
Centre				Candidate			
Centre Number				Number			

#### **READ THESE INSTRUCTIONS FIRST**

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

#### Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

You may use a calculator.

The number of marks is given in brackets [ ] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

DO NOT WRITE IN THE BARCODE.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

For Exam	iner's Use
1	
2	
3	
4	
5	
Total	

## Answer **all** the questions in the spaces provided.

1

١	For
	For Examiner's
ı	Hea

		alt is a pale green crystalline solid which is soluble in water. Mohr's salt is a 'double ch contains
		two cations, one of which is Fe <sup>2+</sup> ,
		one anion which is $SO_4^{2-}$ ,
		and water of crystallisation.
(a)	Soli	e identity of the second cation was determined by the following test.  d Mohr's salt was heated with solid sodium hydroxide and a colourless gas was lived. The gas readily dissolved in water giving an alkaline solution.
	(i)	What is the gas?
	(ii)	What is the formula of the second cation identified by this test?
	(iii)	In this test, a grey/green solid residue was also formed.
		Suggest a name <b>or</b> formula for this solid.
		[3]
(b)	hyd	e identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's . A white precipitate was formed.
	Sug	gest the identity of the white precipitate.
		[1]
(c)		en a double salt such as Mohr's salt is made, the two individual salts are mixed ether in a 1:1 molar ratio, dissolved in water and the solution crystallised.
	(i)	Give the formula of <b>each</b> of the two salts that would be mixed to make the double salt, Mohr's salt.
		salt 1
		salt 2

© UCLES 2006 9701/02/M/J/06

(ii)	Calculate the relative formula mass of <b>each</b> of the salts present in Mohr's salt. salt 1	For Examiner's Use
	relative formula mass of salt 1salt 2	
	relative formula mass of salt 2	
(iii)	The crystals of the double salt contain water of crystallisation.	
	The relative formula mass of Mohr's salt is 392. Use your answers to (ii) to calculate the number of moles of water of crystallisation present in one mole of Mohr's salt.	
	[6]	
	[Total: 10]	

2 The unsaturated hydrocarbon ethyne (acetylene),  $C_2H_2$ , is widely used in 'oxy-acetylene torches' for cutting and welding metals. In the torch, ethyne is burned in oxygen to produce a flame with a temperature of 3400 K.

For Examiner's Use

(a) Ethyne is a linear molecule with a triple bond, C≡C, between the two carbon atoms.

Draw a 'dot-and-cross' diagram of an ethyne molecule.

[1]

**(b)** When used for cutting or welding, ethyne is transported in cylinders which contain the gas under pressure. A typical cylinder has a volume of 76 dm<sup>3</sup> and contains ethyne gas at 1515 kPa pressure at a temperature of 25 °C.

Use the general gas equation, pV = nRT, to calculate the amount, in moles, of ethyne in this cylinder.

[2]

(c) In some countries, ethyne is manufactured from calcium carbide, CaC<sub>2</sub>, which is produced by heating quicklime and coke together at 2300 K.

$$CaO + 3C \rightarrow CaC_2 + CO$$

When water is added to the  $CaC_2$ , calcium hydroxide,  $Ca(OH)_2$ , and ethyne,  $C_2H_2$ , are produced.

(i) Construct a balanced equation for the formation of ethyne from calcium carbide.

(ii) Use this equation and your answer to part (b) to calculate the mass of CaC<sub>2</sub> which will react with an excess of water to produce enough ethyne to fill 100 cylinders of the gas.

[3]

(d)	The equation for the complete combustion of ethyne is given below.
	Use appropriate bond energy data from the <i>Data Booklet</i> to calculate a value for the enthalpy change of combustion of ethyne.
	enthalpy change of combustion of entities.

$$C_2H_2(g) \ + \ {}^5\!\!/_2O_2(g) \ \longrightarrow \ 2CO_2(g) \ + \ H_2O(g)$$

			[3]
(e)	The	e value for the standard enthalpy change of combustion of ethyne is -1300 kJ mol	-1.
	(i)	Define the term standard enthalpy change of combustion.	
	(ii)	Explain why your answer to <b>(d)</b> does not have the same value as the standard enthalpy change of combustion.	ard
			[3]
		[Total:	12]

6 3 This question is about the elements of Group VII, the halogens. (a) Complete the following table. physical state at halogen colour room temperature chlorine bromine iodine [2] (b) Concentrated sulphuric acid is added to separate solid samples of magnesium chloride, magnesium bromide, and magnesium iodide. (i) Describe, in **each** case, **one** observation you would be able to make.  $\mathsf{MgC}l_2$  ..... MgBr<sub>2</sub> ..... Give an equation for the reaction of concentrated sulphuric acid with magnesium chloride. [4] (c) When dilute nitric acid and aqueous silver nitrate are added to a solution of a magnesium halide,  $MgX_2$ , a pale cream precipitate is formed. This precipitate is soluble in concentrated aqueous ammonia but not soluble in dilute aqueous ammonia. What is the identity of the precipitate? Give an equation, with state symbols, for the reaction of the precipitate with

For Examiner's Use

[3]

© UCLES 2006 9701/02/M/J/06

concentrated aqueous ammonia.

(d)		ot glass rod is plunged into separate gas jars, one containing hydrogen chloride and containing hydrogen iodide.
	(i)	For <b>each</b> gas, state what you would observe, if anything, and write an equation for any reaction that takes place.
		HC <i>l</i>
		HI
	(ii)	Explain your answer to (i) in terms of enthalpy changes.
	(11)	Explain your answer to (i) in terms of entiralpy changes.
	(iii)	What is the role of the hot glass rod in any reaction that occurs?
		[6]
		[Total: 15]

4

	onions a	ds containing the allyl group, $\mathrm{CH_2=CHCH_2^-}$ , have pungent smells and are found and garlic. ol, $\mathrm{CH_2=CHCH_2OH}$ , is a colourless liquid which is soluble in water.
(a)	Allyl a	lcohol behaves as an alkene and as a primary alcohol.
	Give t	he structural formula of the organic compound formed when allyl alcohol is
	(i) r	eacted with Br <sub>2</sub> ,
	<b>(ii)</b> h	eated under reflux with an acidified solution of $\text{Cr}_2\text{O}_7^{2-}$ ions.
(b)	When	[2] allyl alcohol is reacted with ${\rm MnO_2}$ at room temperature, propenal, ${\rm CH_2=CHCHO}$ ned.
	What	type of reaction is this?
		[1]
(c)		
(-)	cataly	alcohol may be converted into propanal, $CH_3CH_2CHO$ , by using a ruthenium(IV) st in water.
(-)	сатагу	alcohol may be converted into propanal, $CH_3CH_2CHO$ , by using a ruthenium(IV)
(-)	The re	alcohol may be converted into propanal, $CH_3CH_2CHO$ , by using a ruthenium(IV) st in water.
	The re	alcohol may be converted into propanal, $CH_3CH_2CHO$ , by using a ruthenium(IV) st in water. $CH_2 = CHCH_2OH \xrightarrow{\text{ruthenium(IV) catalyst}} CH_3CH_2CHO$ exactant and the product are isomers.

© UCLES 2006 9701/02/M/J/06

(d)	Allyl alcohol can be converted into propanal in two steps <b>without</b> the use of a ruthenium(IV) catalyst.
	$CH_2 \!\!=\!\! CHCH_2OH  \xrightarrow{step  I}  CH_3CH_2CH_2OH  \xrightarrow{step  II}  CH_3CH_2CHO$
	What reagents and conditions would be used for <b>each</b> step?
	step I
	reagent(s)
	condition(s)
	step II
	reagent(s)
	condition(s)[4]
(e)	By considering your answers to <b>(b)</b> and <b>(d)</b> , suggest what is unusual about the single-step reaction in <b>(c)</b> .
	[1]
(f)	Suggest the structural formula of the organic compound formed when allyl alcohol is
	(i) reacted with cold, dilute MnO <sub>4</sub> <sup>-</sup> ions,
	(ii) heated under reflux with acidified $MnO_4^-$ ions.
	[3]
	[5] [Total: 12]
	[10:α:. 12]

5

		ganic reactions are substitution reactions in which the number of carbon atoms in nic compound is unchanged.
(a)	Wh	at is meant by the term substitution reaction?
		[1]
(b)		e example of a substitution reaction is the formation of an alcohol from a ogenoalkane.
	(i)	Write a balanced equation for the formation of ethanol from bromoethane.
	(ii)	State the conditions for this reaction.
		[2]
(c)		few organic reactions, the product contains <b>one</b> more carbon atom than the starting rerial.
	(i)	Write the equation for a reaction in which the organic compound bromoethane, which contains <b>two</b> carbon atoms, is converted into an organic compound which contains <b>three</b> carbon atoms.
	(ii)	State the conditions for this reaction.
		[2]

© UCLES 2006 9701/02/M/J/06

as the only organic compound.

(d) Ethanol may be converted into propanoic acid in a three-stage process which uses ethanol

	$C_2H_5OH \xrightarrow{\text{step I}} \mathbf{K} \xrightarrow{\text{step II}} \mathbf{L} \xrightarrow{\text{step III}} C_2H_5CO_2H$
(i)	Give the structural formulae of the intermediate compounds ${\bf K}$ and ${\bf L}$ .
	K
	L
(ii)	State the reagent(s) used and give the essential condition(s) for step I and for step III.
	step I
	reagent(s)
	condition(s)
	step III
	reagent(s)
	condition(s)

[Total: 11]

[6]

### **BLANK PAGE**