

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

371397197

CHEMISTRY 9701/02

Paper 2 Structured Questions AS Core

May/June 2007

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

For Examiner's Use				
1				
2				
3				
4				
Total				

This document consists of 9 printed pages and 3 blank pages.



Answer **all** the questions in the spaces provided.

For Examiner's Use

Ethene, C_2H_4 , and hydrazine, N_2H_4 , are hydrides of elements which are adjacent in the Periodic Table. Data about ethene and hydrazine are given in the table below.

	C ₂ H ₄	N_2H_4
melting point/°C	-169	+2
boiling point/°C	-104	+114
solubility in water	insoluble	high
solubility in ethanol	high	high

(a)		ene and hydrazine have a similar arrangement of atoms but differently shaped ecules.	
	(i)	What is the H-C-H bond angle in ethene?	
	(ii)	Draw a 'dot-and-cross' diagram for hydrazine.	
	(iii)	What is the H-N-H bond angle in hydrazine?	
		[4]	
(b)	Sug	melting and boiling points of hydrazine are much higher than those of ethene. gest reasons for these differences in terms of the intermolecular forces each pound possesses.	

© UCLES 2007 9701/02/M/J/07

For Examiner's Use

(c)	-	lain, with the aid of a diagram showing lone pairs of electrons and dipoles, why razine is very soluble in ethanol.
		re)
Eth	ene a	and hydrazine each react with HC l .
		en ethene is reacted with HC l , C $_2$ H $_5$ C l is the only product.
	(i)	Using structural formulae, give an equation for the reaction between ethene and HC1.
	(ii)	What type of reaction occurs between HCl and ethene?
	(iii)	Explain why there is no further reaction between $\mathrm{C_2H_5C}l$ and $\mathrm{HC}l$.
		[3]
(e)	Who	en aqueous hydrazine is reacted with HC l , a solid compound of formula N $_2$ H $_5$ C l may solated. When an excess of HC l is used, a second solid, N $_2$ H $_6$ C l_2 , is formed.
	(i)	Suggest what type of reaction occurs between hydrazine and HCl.
	(ii)	What feature of the hydrazine molecule enables this reaction to occur?
	(iii)	Suggest why one molecule of hydrazine is able to react with one or two molecules of HCl.
		[3]
		[Total: 16]

4 Alcohols and esters are important organic compounds which are widely used as solvents. Esters such as ethyl ethanoate can be formed by reacting carboxylic acids with alcohols. $CH_3CO_2H + C_2H_5OH \rightleftharpoons CH_3CO_2C_2H_5 + H_2O$ This reaction is an example of a dynamic equilibrium. (a) Explain what is meant by the term dynamic equilibrium. **(b)** Write the expression for the equilibrium constant for this reaction, K_c . [1] (c) For this equilibrium, the value of K_c is 4.0 at 298 K. A mixture containing 0.5 mol of ethanoic acid, 0.5 mol ethanol, 0.1 mol ethyl ethanoate and 0.1 mol water was set up and allowed to come to equilibrium at 298 K. The final volume of solution was V dm3. Calculate the amount, in moles, of each substance present at equilibrium. [4]

For Examiner's Use

2

Alcohols may be classified into primary, secondary and tertiary. Some reactions are common to all three types of alcohol. In other cases, the same reagent gives different products depending on the nature of the alcohol.

For Examiner's Use

(d) In the empty squares below give the structural formula of the organic compound formed in each of the reactions indicated.

If no reaction occurs, write 'no reaction' in the space.

alcohol reagent(s) and conditions	CH ₃ CH ₂ CH ₂ CH ₂ OH	CH ₃ CH ₂ CH(OH)CH ₃	(CH ₃) ₃ COH
red phosphorus and iodine heat under reflux			
concentrated H ₂ SO ₄ heat			
Cr ₂ O ₇ ²⁻ /H ⁺ heat under reflux			

[5]

[Total: 11]

For Examiner's Use

- 3 This question is about the elements in Group II of the Periodic Table, magnesium to barium.
 - (a) Complete the table below to show the electronic configuration of calcium atoms and of strontium ions, Sr²⁺.

	1s	2s	2p	3s	Зр	3d	4s	4p	4d
Ca	2	2	6						
Sr ²⁺	2	2	6						

(b)	Ехр	lain the following observations.	[2]
	(i)	The atomic radii of Group II elements increase down the Group.	
	(ii)	The strontium ion is smaller than the strontium atom.	
	(iii)	The first ionisation energies of the elements of Group II decrease with increas proton number.	ing
			 [4]

(c)	som	nples of magnesium and calcium are placed separately in cold water and left for ne time. In each case , describe what you would see and write a balanced equation each reaction.	For Examiner's Use
	(i)	magnesium	
		observation	
		equation	
	(ii)	calcium	
		observation	
		equation[6]	
		[O]	
(d)	Stro	ontium nitrate, Sr(NO ₃) ₂ undergoes thermal decomposition.	
	(i)	State one observation you would make during this reaction.	
	(ii)	Write a balanced equation for this reaction.	
		[4]	
		[Total: 16]	
			1

4		emmercial paint and varnish removers contain a mixture of dichloromethane, ${\rm CH_2Cl_2}$, and ethanol, ${\rm CH_3OH}$.						
	(a)	In e	at would be observed when the following reactions are carried out? ach case, give the name or formula of the reaction product which is responsible for observation you have made.					
		(i)	$\mathrm{CH_2Cl_2}$ is reacted with NaOH(aq) and $\mathrm{AgNO_3}(\mathrm{aq})$ and the mixture left to stand.					
			observation					
			product responsible					
		(ii)	CH ₃ OH is mixed with PCl ₅ .					
			observation					
			product responsible					
		(iii)	CH ₃ OH is reacted with sodium.					
			observation					
			product responsible					
			[6]					
	(b)	Who forn	en $\mathrm{CH_2Cl_2}$ is heated under reflux with an excess of NaOH(aq), a compound \mathbf{W} is ned.					
		W h	as the following composition by mass: C, 40.0%; H, 6.7%; O, 53.3%.					
		Use CH ₂	this information and the $\textit{Data Booklet}$ to show that the empirical formula of \mathbf{W} is ${}_{2}^{O}$ O.					
			[2]					

For Examiner's Use

			9					
(c)	Con	npounds with the empiri	cal formula CH ₂ O can h	nave the molecular form	ula C ₂ H ₄ O ₂ .			
		possible structural for D ₂ CH ₃ and H ₂ C=C(OH)		with molecular formula	C ₂ H ₄ O ₂ are			
		ne boxes below, draw d molecular formula C ₂ H ₄		hree further structural	isomers with			
	Do not attempt to draw any structures containing rings or O-O bonds.							
		x	Υ	Z				
(d)	d) Identify which of your compounds, X , Y , or Z , will react with the following reagents. In each case, state what you would observe.							
	(i)	solid NaHCO ₃						
		compound						
		observation						
	(ii)	Tollens' reagent						
		compound						
		observation			[4]			
(e)	One	e of the three compound	ls, X , Y , or Z , shows ste	ereoisomerism.	[+]			
	Draw displayed, labelled structures of the stereoisomers of this compound.							

[2]

For Examiner's Use

[Total: 17]

© UCLES 2007 9701/02/M/J/07

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.